


Product datasheet

Anti-IRAK1 antibody ab85071

★ ★ ★ ☆ ☆ 1 Abreviews 3 Images

Overview

Product name	Anti-IRAK antibody
Description	Rabbit polyclonal to IRAK
Host species	Rabbit
Tested applications	Suitable for: WB, ICC/IF
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat, Cow, Macaque monkey 
Immunogen	Synthetic peptide conjugated to KLH derived from within residues 350 - 450 of Human IRAK. Read Abcam's proprietary immunogen policy (Peptide available as ab99364 .)
Positive control	<div style="border: 1px solid #ccc; padding: 5px; display: inline-block;"> Purchase matching WB positive control: Recombinant Human IRAK protein > </div> <p>This antibody gave a positive signal in Human liver tissue lysate.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab85071** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★☆	Use a concentration of 1 µg/ml. Detects a band of approximately 74 kDa (predicted molecular weight: 76 kDa).
ICC/IF		Use a concentration of 5 µg/ml.

Target

Function

Binds to the IL-1 type I receptor following IL-1 engagement, triggering intracellular signaling cascades leading to transcriptional up-regulation and mRNA stabilization. Isoform 1 binds rapidly but is then degraded allowing isoform 2 to mediate a slower, more sustained response to the cytokine. Isoform 2 is inactive suggesting that the kinase activity of this enzyme is not required for IL-1 signaling. Once phosphorylated, IRAK1 recruits the adapter protein PELI1.

Tissue specificity

Isoform 1 and isoform 2 are ubiquitously expressed in all tissues examined, with isoform 1 being more strongly expressed than isoform 2.


Sequence similarities

Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. Pelle subfamily. Contains 1 protein kinase domain.

Post-translational modifications

Autophosphorylated or is transphosphorylated by IRAK4 following recruitment to the IL-1RI. In the case of isoform 1, this is linked to ubiquitination and degradation. Polyubiquitinated; after cell stimulation with IL-1-beta. Polyubiquitination occurs with polyubiquitin chains linked through 'Lys-63'.

Images



Lane 1: Wild type HAP1 whole cell lysate (20 μ g)


Lane 2: IRAK1 knockout HAP1 whole cell lysate (20 μ g)

Lane 3: HeLa whole cell lysate (20 μ g)

Lane 4: Hek293 whole cell lysate (20 μ g)

Lanes 1 - 4: Merged signal (red and green). Green - ab85071 observed at 85 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab85071 was shown not to specifically react with IRAK1 when IRAK1 knockout samples were used. Wild-type and IRAK1 knockout samples were subjected to SDS-PAGE. Ab85071 and [ab8245](#) (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1 μ g/ml and 1/10000 dilution respectively. Blots were developed with 800CW Goat anti Rabbit and 680CW Goat anti Mouse secondary antibodies at 1/10012 dilution for 1 hour at room temperature before imaging.



Western blot - IRAK antibody (ab85071)

Anti-IRAK1 antibody (ab85071) at 1 µg/ml +
Human liver tissue lysate - total protein
(ab29889) at 10 µg

Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-
Adsorbed (HRP) at 1/3000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 76 kDa

Observed band size: 74 kDa

Additional bands at: 36 kDa. We are unsure
as to the identity of these extra bands.

Exposure time: 3 minutes



Immunocytochemistry/ Immunofluorescence - IRAK
antibody (ab85071)

ICC/IF image of ab85071 stained MCF-7
cells. The cells were 4% formaldehyde fixed
(10 min) and then incubated in 1%BSA / 10%
normal goat serum / 0.3M glycine in 0.1%
PBS-Tween for 1h to permeabilise the cells
and block non-specific protein-protein
interactions. The cells were then incubated
with the antibody ab85071 at 5µg/ml overnight
at +4°C. The secondary antibody (green) was
Alexa Fluor® 488 goat anti- rabbit IgG (H+L)
used at a 1/1000 dilution for 1h. Alexa Fluor®
594 WGA was used to label plasma
membranes (red) at a 1/200 dilution for 1h.
DAPI was used to stain the cell nuclei (blue) at
a concentration of 1.43µM. This antibody also
gave a positive result in Methanol fixed (100%,
5min) MCF-7 cells at 5ug/ml

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors