


Product datasheet

Anti-TRIM23 antibody ab90282

1 Image

Overview

| | |
|----------------------------|--|
| Product name | Anti-TRIM23 antibody |
| Description | Rabbit polyclonal to TRIM23 |
| Host species | Rabbit |
| Tested applications | Suitable for: WB |
| Species reactivity | Reacts with: Human Predicted to work with: Mouse, Rat, Rabbit, Horse, Chicken, Guinea pig, Cow, Cat, Dog, Zebrafish  |
| Immunogen | Synthetic peptide from a region within internal sequence amino acids 216-265 (HSVLEPEANQ IRASILDMAH CIRTFTEEIS DYSRKLVGV QHIEGGEQV) of Human TRIM23 (NP_150231) |
| Positive control | Human fetal spleen lysate |

Properties

| | |
|-----------------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |
| Storage buffer | Preservative: None Constituents: 2% Sucrose, PBS |
| Purity | Immunogen affinity purified |
| Clonality | Polyclonal |
| Isotype | IgG |

Applications

Our [Abpromise guarantee](#) covers the use of **ab90282** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| WB | | Use a concentration of 1 µg/ml. Predicted molecular weight: 64 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T. |

Target

| | |
|------------------------------|--|
| Function | Acts as an E3 ubiquitin-protein ligase. In the presence of the human cytomegalovirus (HCMV) protein UL144, participates in 'Lys-63'-linked auto-ubiquitination of TRAF6 resulting in the virally controlled activation of NF-kappa-B at early time of infection. The C-terminus can act as an allosteric activator of the cholera toxin catalytic subunit. |
| Pathway | Protein modification; protein ubiquitination. |
| Sequence similarities | In the C-terminal section; belongs to the small GTPase superfamily. Arf family. Contains 1 B box-type zinc finger. Contains 1 RING-type zinc finger. |
| Domain | The RING-type zinc finger domain is responsible for E3 ubiquitin ligase activity. |
| Cellular localization | Endomembrane system. Golgi apparatus membrane. Lysosome membrane. Membrane-associated with the Golgi complex and lysosomal structures. |

Images



Anti-TRIM23 antibody (ab90282) at 1 µg/ml +
Human fetal spleen lysate at 10 µg

Secondary

HRP conjugated anti-Rabbit IgG at 1/50000
dilution

Predicted band size: 64 kDa

Observed band size: 64 kDa

Gel concentration: 12%

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors