

Product datasheet

Anti-Myc tag antibody [9E10] (FITC) ab117599

1 References

Overview

Product name	Anti-Myc tag antibody [9E10] (FITC)
Description	Mouse monoclonal [9E10] to Myc tag (FITC)
Conjugation	FITC. Ex: 493nm, Em: 528nm
Specificity	This antibody is specific for Myc tagged proteins. The Myc tag epitope (EQKLISEEDL) is located at the dimerization site of c-myc and therefore this antibody does not perform well at recognizing endogenous c-myc.
Tested applications	Suitable for: Flow Cyt
Immunogen	Synthetic peptide: AEEQKLISEEDLL , corresponding to C terminal amino acids 408 - 420 of Human c-Myc
	Run BLAST with Run BLAST with

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.09% Sodium azide Constituents: 1% BSA, 98% PBS
Purity	Protein G purified
Clonality	Monoclonal
Clone number	9E10
Isotype	IgG1

Applications

Our [Abpromise guarantee](#) covers the use of **ab117599** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		1/10. Membrane permeabilisation is required for this application. Use 10µl of antibody to label 10 ⁶ cells in 100µl. ab91356 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Target

Relevance

Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells.

Cellular localization

Nuclear

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors