

Product datasheet

Anti-cetA antibody ab22172

1 Image

Overview

Product name	Anti-cetA antibody
Description	Mouse polyclonal to cetA
Tested applications	Suitable for: WB
Species reactivity	Reacts with <i>Campylobacter jejuni</i> . Not yet tested in other species.
Immunogen	Fusion protein: SVVKIDHILYKSNMNLNNGAQNFNLESVDPISNLCQDERAQQVINELSS ETELNLAKEFIKDNAKKAIEESSQDYDQKAYDAVNDIKSLEQRSAIL , corresponding to amino acids 355/454 of <i>Campylobacter jejuni</i> methyl-accepting chemotaxis protein.
	Run BLAST with Run BLAST with

General notes Produced from outbred CD1 mice

This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang *et al.* [PubMed: 1545867](#); Chambers and Johnston [PubMed: 12910245](#); Barry and Johnston [PubMed: 9234514](#)). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an *E.coli* lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
Storage buffer	Constituents: 50% Glycerol
Purity	Whole antiserum

Primary antibody notes

This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang *et al.* [PubMed: 1545867](#); Chambers and Johnston [PubMed: 12910245](#); Barry and Johnston [PubMed: 9234514](#)). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an *E.coli* lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

Clonality

Polyclonal

Isotype

IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab22172** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		1/1000. Predicted molecular weight: 51 kDa. This antibody has been tested in Western blot against an <i>E.coli</i> lysate containing the partial recombinant fusion protein used as an immunogen. We have no data on detection of endogenous protein.

Target

Relevance

Bacterial chemotactic-signal transducers are proteins that respond to changes in the concentration of attractants and repellents in the environment, and transduce a signal from the outside to the inside of the cell. These proteins undergo two covalent modifications: deamidation and reversible methylation. Attractants increase the level of methylation while repellents decrease it. The methyl groups are added by the methyl-transferase cheR and are removed by the methylesterase cheB

Images



All lanes : Anti-cetA antibody (ab22172) at 1/1000 dilution

Lane 1 : Total protein extract from E. coli with ~50ng to 100ng of a negative control fusion protein with an irrelevant antigen at 20 ug

Lane 2 : Total protein extract from E. coli with ~50ng to 500ng of the antigen fusion protein at 20 ug

Secondary

Rabbit anti-mouse IgG + IgM, (H+L) horseradish peroxidase conjugated at 1/5000 dilution

Predicted band size : 51 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors