abcam

Product datasheet

Dienogest, Steroidal progesterone ab141282

1 References 2 Images

Overview

Product name Dienogest, Steroidal progesterone

Description Steroidal progesterone

Biological descriptionSteroidal progesterone. Non-ethinylated progestin structurally related to testosterone. Anti-

androgenic and antiproliferative effects in vitro.

Purity > 99%

CAS Number 65928-58-7

Chemical structure

CH OH

Properties

Chemical name (17α)-17-Hydroxy-3-oxo-19-norpregna-4,9-diene-21-nitrile

Molecular weight 311.42

PubChem identifier 68861

Storage instructions Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12

months.

Solubility overview Soluble in DMSO to 100 mM

Handling Wherever possible, you should prepare and use solutions on the same day. However, if you need

to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and

prior to opening the vial we recommend that you allow your product to equilibrate to room $\frac{1}{2} \int_{\mathbb{R}^{n}} \left(\frac{1}{2} \int$

temperature for at least 1 hour.

Toxic, refer to SDS for further information

Need more advice on solubility, usage and handling? Please visit our frequently asked

questions (FAQ) page for more details.

SMILES O=C3C=C4CC[C@@H]2C(CC[C@@]1(C)[C@H]2CC[C@@]1(O)CC#N)=C4CC3

1


Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab141282 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
Functional Studies		Use at an assay dependent concentration.

Images



Chemical Structure - Dienogest, Steroidal progesterone (ab141282)

2D chemical structure image of ab141282, Dienogest, Steroidal progesterone



Functional Studies - Dienogest, Steroidal progesterone (ab141282)

<u>ab16895</u> staining MDM2 in MCF7 cells treated with dienogest (ab141282), by ICC/IF. Decrease in MDM2 expression correlates with increased concentration of dienogest, as described in literature.

The cells were incubated at 37° C for 24 hour in media containing different concentrations of ab141282 (dienogest) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab16895 (5 µg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 anti-mouse polyclonal antibody (ab96879) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

- · Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team