

Product datasheet

10x Citrate Buffer pH 6.0 ab64214

★★★★★ [1 Abreviews](#) [8 References](#) [2 Images](#)

Overview

Product name	10x Citrate Buffer pH 6.0
Tested applications	<b>Suitable for:</b> IHC-P
General notes	Citrate buffer pH 6.0 for heat-induced antigen retrieval (HIER) during IHC.

Properties


Form	Liquid
Storage instructions	Shipped at 4°C. Shipped at Room Temperature. Store at Room Temperature.
Storage buffer	pH: 6.00 Constituent: Citrate Buffer  Buffer x10 concentrated.
Relevance	Citrate buffer is used on formalin-fixed, paraffin-embedded tissue sections mounted on glass slides for target retrieval prior to immunohistochemistry (IHC) procedures. Target retrieval prior to IHC procedures obtains positive results, or results in an increase in staining intensity with many primary antibodies.

Applications

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab64214 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent dilution.

Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - 10x Citrate Buffer pH 6.0 (ab64214)

Juho T et al. PLoS One (2014) Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>


Immunohistochemical analysis of mouse lung. Samples were fixed in 10% buffered formalin, paraffinized and sliced at 1.5  $\mu$ m thick. Antigen retrieval was performed using ab64214 for the deparaffinized slices. Sections were blocked with 2% normal goat serum, PBS(-) and 0.1% Tween20. They were then incubated with the primary antibodies for 1 hour at 4°C and with secondary antibodies for 30 minutes at room temperature. The avidin-biotin-peroxidase complex method with peroxidase streptavidin and the DAB substrate kit **ab64238** was performed.

(A) A resected lung from a mouse sacrificed 28 days after SCL injection.

(B) Haematoxylin and Eosin staining showing the tumor composed of a central area with necrosis and a peripheral zone filled with SCLs.

(C) Immunohistochemical staining for MMP-14 showing positive expression of MMP-14 in the peripheral zone of the tumor and negative in central zone.

(D) Immunohistochemical staining weak staining of MMP-2 in the peripheral zone.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - 10x Citrate Buffer pH 6.0 (ab64214)

Juho T et al. PLoS One (2014)

Immunohistochemical analysis of liver tissue section labeling PCNA with **ab18197** at 1/250. Antigen retrieval was performed with antigen retrieval buffer, **ab64236**. A and B, PCNA staining in right and left grafts 1 day after reperfusion. E and F, PCNA staining in right and left grafts 5 day after reperfusion.

This image was produced using **ab64236** (a higher concentration of ab64214 ). Therefore, we expect ab64214 to work as well when diluted to the required concentration.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors