

SW480 whole cell lysate ab3957

1 References

Overview

Product name	SW480 whole cell lysate
General notes	<p>Cell line: SW480 (Human colorectal adenocarcinoma).</p> <p>Growth media: Leibovitz's L-15 medium with 2 mM L-glutamine, and 10% FBS.</p> <p>SW480 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylene diamine tetra acetic acid, 1 mM phenyl methyl sulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecyl sulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (0.045 M Tris-HCl pH 6.8, 10% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue), containing 0.05 M DTT.</p>
Tested applications	Suitable for: WB

Properties

Mycoplasma free	Yes
Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 7.20</p> <p>Constituent: 100% SDS Sample Buffer</p>
Lysate notes	<p>SW480 cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylene diamine tetra acetic acid, 1 mM phenyl methyl sulfonyl flouride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecyl sulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecyl sulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.</p>
Background	Cells are epithelial like in morphology and used as an in vitro model for colorectal cancer to study tumor markers, CEA production, biochemistry of tumorigenicity, pharmacodynamics, hormone sensibility and as a positive control for the nuclear protein Beta catenin.

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab3957 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. SW480 cell lysate is ready to load on SDS-PAGE for Western blotting, 10 - 20 µg per lane is recommended for mini gel.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors