abcam

Product datasheet

Anti-AACT antibody [EPR14118(B)] ab184567

Recombinant RabMAb

1 References 2 Images

Overview

Product name Anti-AACT antibody [EPR14118(B)]

Description Rabbit monoclonal [EPR14118(B)] to AACT

Host species Rabbit

Suitable for: WB **Tested applications**

Species reactivity Reacts with: Human

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

Positive control Human serum and Human plasma.

This product is a recombinant monoclonal antibody, which offers several advantages including: **General notes**

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

Purity Tissue culture supernatant

Clonality Monoclonal Clone number EPR14118(B)

Isotype lgG

Applications

The Abpromise guarantee

Our <u>Abpromise guarantee</u> covers the use of ab184567 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000 - 1/10000. Detects a band of approximately 70 kDa (predicted molecular weight: 48 kDa).


Target

Function	Although its physiological function is unclear, it can inhibit neutrophil cathepsin G and mast cell chymase, both of which can convert angiotensin-1 to the active angiotensin-2.	
Tissue specificity	Plasma. Synthesized in the liver. Like the related alpha-1-antitrypsin, its concentration increases in the acute phase of inflammation or infection. Found in the amyloid plaques from the hippocampus of Alzheimer disease brains.	
Involvement in disease	Defects in SERPINA3 may be a cause of chronic obstructive pulmonary disease (COPD) [MIM:107280].	
Sequence similarities	Belongs to the serpin family.	
Domain	The reactive center loop (RCL) extends out from the body of the protein and directs binding to the target protease. The protease cleaves the serpin at the reactive site within the RCL, establishing a covalent linkage between the carboxyl group of the serpin reactive site and the serine hydroxyl of the protease. The resulting inactive serpin-protease complex is highly stable.	

Images

(ab184567)

Cellular localization



Secondary
All lanes : A

All lanes: Anti-Rabbit IgG (HRP), specific to the non-reduced form

of IgG at 1/1000 dilution

Predicted band size: 48 kDa


All lanes : Anti-AACT antibody [EPR14118(B)] (ab184567) at

1/10000 dilution

Secreted.

Lane 1 : Human serum
Lane 2 : Human plasma

Lysates/proteins at 10 µg per lane.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors