abcam

Product datasheet

Anti-Actin antibody [MAC 237] ab50591

★★★★★ 1 Abreviews 9 References 2 Images

Overview

Product name Anti-Actin antibody [MAC 237]

Description Rat monoclonal [MAC 237] to Actin

Host species Rat

Specificity Antibody reacts with actin in Lethocerus and Drosophila flight and non-flight muscle and with actin

in flight muscles of all other insect species tested. Also reacts with mammalian actin (tested vs.

rat). Antibody reacts with arthrin in flight muscles of Hemiptera and Diptera.

Tested applications Suitable for: IHC-P

Species reactivity Reacts with: Human

Immunogen Tissue, cells or virus corresponding to Waterbug Actin. Flight muscle extract from Lethocerus

indicus (Waterbug)

Positive controlThis antibody gave a positive result in IHC in the following FFPE tissue: Human normal skin.

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

Storage buffer Preservative: 0.1% Sodium azide

Constituent: PBS

Purity Protein A purified

Clonality Monoclonal

Clone number MAC 237

Isotype IgG2b

1

Applications

The Abpromise quarantee

Our Abpromise guarantee covers the use of ab50591 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	****(1)	Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Target

Function

Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

Involvement in disease

Defects in ACTA1 are the cause of nemaline myopathy type 3 (NEM3) [MIM:161800]. A form of nemaline myopathy. Nemaline myopathies are muscular disorders characterized by muscle weakness of varying severity and onset, and abnormal thread-or rod-like structures in muscle fibers on histologic examination. The phenotype at histological level is variable. Some patients present areas devoid of oxidative activity containg (cores) within myofibers. Core lesions are unstructured and poorly circumscribed.

Defects in ACTA1 are a cause of myopathy congenital with excess of thin myofilaments (MPCETM) [MIM:161800]. A congenital muscular disorder characterized at histological level by areas of sarcoplasm devoid of normal myofibrils and mitochondria, and replaced with dense masses of thin filaments. Central cores, rods, ragged red fibers, and necrosis are absent. Defects in ACTA1 are a cause of congenital myopathy with fiber-type disproportion (CFTD) [MIM:255310]; also known as congenital fiber-type disproportion myopathy (CFTDM). CFTD is a genetically heterogeneous disorder in which there is relative hypotrophy of type 1 muscle fibers compared to type 2 fibers on skeletal muscle biopsy. However, these findings are not specific and can be found in many different myopathic and neuropathic conditions.


Sequence similarities

Cellular localization

Belongs to the actin family.

Cytoplasm > cytoskeleton.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Actin antibody [MAC 237] (ab50591)

This image is courtesy of an Abreview submitted by Anne Sailer

ab50591 staining Actin in mouse proximal colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffinembedded sections). Tissue was fixed with formaldehyde and blocked with 5% NDS for 24 hours at 4°C; antigen retrieval was by heat mediation in buffer, pH9. Samples were incubated with primary antibody (1/100 in 5% BSA/NDS) for 24 hours at 4°C. An undiluted Alexa Fluor[®] 647-conjugated donkey anti-rat IgG polyclonal was used as the secondary antibody.

Also stained with ab76055 (green) at 1/100.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Actin antibody [MAC 237] (ab50591)

IHC image of Actin staining in Human normal skin formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol B. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab50591, 5µg/ml, for 15 mins at room temperature. A Goat anti-Rat biotinylated secondary antibody was used to detect the primary, and visualized using an HRP conjugated ABC system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors