

## Product datasheet

# Agarose Anti-DDDDK tag (Binds to FLAG® tag sequence) antibody ab1240

## 5 References

### Overview

<b>Product name</b>	Agarose Anti-DDDDK tag (Binds to FLAG® tag sequence) antibody
<b>Description</b>	Agarose Goat polyclonal to DDDDK tag (Binds to FLAG® tag sequence)
<b>Host species</b>	Goat
<b>Conjugation</b>	Agarose
<b>Tested applications</b>	<b>Suitable for:</b> IP
<b>Species reactivity</b>	<b>Reacts with:</b> Species independent
<b>Immunogen</b>	Full length native protein (purified) corresponding to DDDDK tag conjugated to Keyhole Limpet Haemocyanin (KLH). Sequence: xxxDDDDK

 [Run BLAST with](#)

 [Run BLAST with](#)

### General notes

Epitope tags provide a method to localize gene products in a variety of cell types, study the topology of proteins and protein complexes, identify associated proteins, and characterize newly identified, low abundance or poorly immunogenic proteins when protein specific antibodies are not available. Tagging with xxxDDDDK may be done at the N-terminus, N-terminus preceded by a methionine residue, C-terminus, and in internal positions of the target protein. The small size of the epitope tag and its high hydrophilicity tend to decrease the possibility of interference with protein expression, proteolytic maturation, antigenicity and function. The enterokinase cleavage site allows it to be completely removed from the purified fusion proteins.

Affinity purified antibodies were coupled to agarose beads using a cyanogen bromide method.

FLAG® is a registered trade mark of Sigma Aldrich Biotechnology LP. It is used here for informational purposes only.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

## Properties

---

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	pH: 6.8 Preservative: 0.1% Sodium azide Constituents: 0.0268% PBS, 0.58% Sodium chloride
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

## Applications

---

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab1240 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IP		Use at an assay dependent dilution.

## Target

---

**Relevance** This is a useful tool for the localisation and characterisation of DDDDK tagged proteins (Binds to FLAG® tag sequence).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors