abcam

Product datasheet

Anti-Alpha Skeletal Muscle Actin antibody ab52218

★★★★★ 1 Abreviews 8 References 3 Images

Overview

Product name Anti-Alpha Skeletal Muscle Actin antibody

Description Rabbit polyclonal to Alpha Skeletal Muscle Actin

Host species Rabbit

Specificity Actin alpha 1 antibody detects endogenous levels of total Actin alpha 1 protein

Tested applications
Suitable for: WB, IHC-P
Species reactivity
Reacts with: Human

Immunogen Synthetic derived from human Actin alpha 1.

General notes

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle.

Storage buffer pH: 7

Preservative: 0.02% Sodium azide

Constituents: PBS, 0.87% Sodium chloride, 50% Glycerol (glycerin, glycerine)

Without Mg+2 and Ca+2

Purity Immunogen affinity purified

Purification notes The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-

specific immunogen.

Clonality Polyclonal

Isotype IgG

1

Applications

The Abpromise guarantee

Our Abpromise quarantee covers the use of ab52218 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/1000. Detects a band of approximately 45 kDa (predicted molecular weight: 42 kDa).
IHC-P	★★★★☆ (1)	Use at an assay dependent concentration.

Target

Function

Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

Involvement in disease

Defects in ACTA1 are the cause of nemaline myopathy type 3 (NEM3) [MIM:161800]. A form of nemaline myopathy. Nemaline myopathies are muscular disorders characterized by muscle weakness of varying severity and onset, and abnormal thread-or rod-like structures in muscle fibers on histologic examination. The phenotype at histological level is variable. Some patients present areas devoid of oxidative activity containg (cores) within myofibers. Core lesions are unstructured and poorly circumscribed.

Defects in ACTA1 are a cause of myopathy congenital with excess of thin myofilaments (MPCETM) [MIM:161800]. A congenital muscular disorder characterized at histological level by areas of sarcoplasm devoid of normal myofibrils and mitochondria, and replaced with dense masses of thin filaments. Central cores, rods, ragged red fibers, and necrosis are absent. Defects in ACTA1 are a cause of congenital myopathy with fiber-type disproportion (CFTD) [MIM:255310]; also known as congenital fiber-type disproportion myopathy (CFTDM). CFTD is a genetically heterogeneous disorder in which there is relative hypotrophy of type 1 muscle fibers compared to type 2 fibers on skeletal muscle biopsy. However, these findings are not specific and can be found in many different myopathic and neuropathic conditions.


Sequence similarities

Cellular localization

Belongs to the actin family.

Cytoplasm > cytoskeleton.

Images



embedded sections) - Anti-Alpha Skeletal Muscle Actin antibody (ab52218)

Immunohistochemistry (Formalin/PFA-fixed paraffin-

All lanes: Anti-Alpha Skeletal Muscle Actin antibody (ab52218) at 1/500 dilution

Lane 1: extracts from LOVO cells,

Lane 2: extracts from LOVO cells, with immunizing peptide

(overnight incubation), and amplification may be required.

Ab52218 staining human skeletal muscle tissue. Staining is

Sections were stained using an automated system DAKO

Left panel: with primary antibody at 2 ug/ml. Right panel: isotype

Autostainer Plus, at room temperature. Sections were rehydrated and antigen retrieved with the Dako 3-in-1 AR buffer EDTA pH 9.0

in a DAKO PT Link. Slides were peroxidase blocked in 3% H2O2 in methanol for 10 minutes. They were then blocked with Dako

Protein block for 10 minutes (containing casein 0.25% in PBS) then


incubated with primary antibody for 20 minutes and detected with

Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time


localised to cytoplasm.

control.

Predicted band size: 42 kDa Observed band size: 45 kDa



Western blot - Anti-Alpha Skeletal Muscle Actin antibody (ab52218)



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Alpha Skeletal Muscle
Actin antibody (ab52218)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human muscle tissue labelling Alpha Skeletal Muscle Actin with ab52218. Right pane - treated with immunizing peptide.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors