

Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free ab210714

KO VALIDATED

Recombinant

RabMAb

10 Images

Overview

Product name	Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free
Description	Rabbit monoclonal [Y365] to AMPK alpha 1 - BSA and Azide free
Host species	Rabbit
Specificity	This antibody is specific for human AMPK alpha 1. This antibody shows low affinity on mouse and rat samples.
Tested applications	Suitable for: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: HeLa, HepG2, C6, NIH/3T3 and MCF-7 cell lysate. Mouse liver, brain, retina, and skeletal muscle tissue lysates. IHC-P: Human cervical carcinoma and lung carcinoma tissues. ICC/IF: MCF-7 cells. Flow Cyt (intra): HeLa cells. IP: HeLa cell lysate.
General notes	ab210714 is the carrier-free version of ab32047 .

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our [conjugation kits](#) for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.20 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	Y365
Isotype	IgG

Applications


The Abpromise guarantee Our [**Abpromise guarantee**](#) covers the use of ab210714 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		Use at an assay dependent concentration. Predicted molecular weight: 63 kDa.
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols .
ICC/IF		Use at an assay dependent concentration.

Target

Function	Responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. It also regulates cholesterol synthesis via phosphorylation and inactivation of hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Appears to act as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-AMP rises in response to fuel limitation and/or hypoxia. This is a catalytic subunit.
Sequence similarities	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. SNF1 subfamily. Contains 1 protein kinase domain.

Images



Western blot - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

All lanes : Anti-AMPK alpha 1 antibody [Y365] ([ab32047](#)) at 1/1000 dilution

Lane 1 : Wild-type RAW 264.7 cell lysate

Lane 2 : PRKAA1 knockout RAW 264.7 cell lysate

Lane 3 : Mouse Liver cell lysate


Lane 4 : Neuro2A cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 63 kDa


False colour image of Western blot: Anti-AMPK alpha 1 antibody [Y365] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab32047](#) was shown to bind specifically to AMPK alpha 1. A band was observed at 64 kDa in wild-type RAW 264.7 cell lysates with no signal observed at this size in PRKAA1 knockout cell line [ab280055](#) (knockout cell lysate [ab280114](#)). To generate this image, wild-type and PRKAA1 knockout RAW 264.7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling AMPK alpha 1 with purified **ab32047** at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).




Immunocytochemistry/ Immunofluorescence - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Immunocytochemistry/Immunofluorescence analysis of MCF7 cells labelling AMPK alpha 1 with purified **ab32047** at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/250) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Intracellular Flow Cytometry analysis of HeLa cells labelling AMPK alpha 1 with purified **ab32047** at 1/150 (red). Cells were fixed with 2% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Immunoprecipitation - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

ab32047 (purified) at 1/40 immunoprecipitating AMPK alpha 1 in HeLa whole cell lysate.

Lane 1 (input): HeLa whole cell lysate (10µg)

Lane 2 (+): **ab32047** + HeLa whole cell lysate (10µg).


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab32047** in HeLa whole cell lysate.

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/1500 dilution.

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Western blot - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

All lanes : Anti-AMPK alpha 1 antibody [Y365] ([ab32047](#)) at 1/500 dilution

Lane 1 : HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

Lane 3 : Mouse liver tissue lysate

Lane 4 : Mouse brain tissue lysate

Lane 5 : Mouse kidney tissue lysate

Lane 6 : Mouse retina tissue lysate

Lane 7 : Mouse skeletal muscle tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 63 kDa


Observed band size: 63 kDa

Additional bands at: 40 kDa (possible non-specific binding)

Exposure time: 3 minutes

Blocking/Diluting buffer and concentration: 5% NFDM/TBST.

This antibody shows low affinity on mouse samples. This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32047](#)).



All lanes : Anti-AMPK alpha 1 antibody [Y365] ([ab32047](#)) at 1/20000 dilution

Lane 1 : MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate

Lane 2 : HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

Lane 3 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 4 : K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

Lane 5 : C6 (Rat glial tumor glial cell) whole cell lysate

Lane 6 : Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysate

Lane 7 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 63 kDa


Observed band size: 63 kDa

Exposure time: 180 seconds

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

This antibody shows low affinity on mouse and rat samples.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32047](#)).



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Overlay histogram showing HeLa cells stained with unpurified **ab32047** (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (unpurified **ab32047**, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervical carcinoma tissue labelling AMPK alpha 1 with unpurified **ab32047** at a dilution of 1/100.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors