abcam

Product datasheet

Anti-ATG7 antibody ab99001

2 References

Overview

Product name Anti-ATG7 antibody

Description Rabbit polyclonal to ATG7

Host species Rabbit

Tested applications Suitable for: WB

Species reactivity Reacts with: Arabidopsis thaliana

Immunogen Full length protein corresponding to ATG7.

General notesThe Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.

Storage buffer Constituent: Whole serum

Purity Whole antiserum

Clonality Polyclonal

Isotype IgG

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab99001 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration.

Target

laiget	
Function	E1-like activating enzyme involved in the 2 ubiquitin-like systems required for cytoplasm to vacuole transport (Cvt) and autophagy. Activates ATG12 for its conjugation with ATG5 as well as the ATG8 family proteins for their conjugation with phosphatidylethanolamine. Both systems are needed for the ATG8 association to Cvt vesicles and autophagosomes membranes. Required for autophagic death induced by caspase-8 inhibition. Required for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Modulates p53/TP53 activity to regulate cell cycle and survival during metabolic stress. Plays also a key role in the maintenance of axonal homeostasis, the prevention of axonal degeneration, the maintenance of hematopoietic stem cells, the formation of Paneth cell granules, as well as in adipose differentiation.
Tissue specificity	Widely expressed, especially in kidney, liver, lymph nodes and bone marrow.
Sequence similarities	Belongs to the ATG7 family.
Domain	The C-terminal part of the protein is essential for the dimerization and interaction with ATG3 and ATG12. The N-terminal FAP motif (residues 15 to 17) is essential for the formation of the ATG89-PE and ATG5-ATG12 conjugates.
Post-translational modifications	Acetylated by EP300.
Cellular localization	Cytoplasm. Preautophagosomal structure. Localizes also to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

•	Guarantee only valid for products bought direct from Abcam or one of our authorized distributors				
		3			