


Product datasheet

Biotin Anti-ICAM1 antibody ab7815

★★★★★ [3 Abreviews](#) [7 References](#) [1 Image](#)

Overview

Product name	Biotin Anti-ICAM1 antibody
Description	Biotin Rabbit polyclonal to ICAM1
Host species	Rabbit
Conjugation	Biotin
Tested applications	Suitable for: Sandwich ELISA
Species reactivity	Reacts with: Human Predicted to work with: Chimpanzee, Baboon, Cynomolgus monkey, Rhesus monkey 
Immunogen	Full length protein corresponding to Human ICAM1. Database link: P05362
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.065% Sodium azide Constituent: 0.1% BSA
Purification notes	0.2 µm filtered
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab7815 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Sandwich ELISA		Use a concentration of 0.5 µg/ml. Can be paired for Sandwich ELISA with Mouse monoclonal [MEM-111] to ICAM1 (ab2213) . For sandwich ELISA, use this antibody as Detection at 0.5µg/ml with ab2213 as Capture.

Target

Function

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHO G activation. In case of rhinovirus infection acts as a cellular receptor for the virus.

Sequence similarities

Belongs to the immunoglobulin superfamily. ICAM family.
Contains 5 Ig-like C2-type (immunoglobulin-like) domains.


Post-translational modifications

Monoubiquitinated, which is promoted by MARCH9 and leads to endocytosis.

Cellular localization

Membrane.

Images



Standard Curve for ICAM1 (Analyte: **ab82125**) dilution range 1pg/ml to 1ug/ml using Capture Antibody Mouse monoclonal [MEM-111] to ICAM1 (**ab2213**) at 5ug/ml and Detector Antibody Rabbit polyclonal to ICAM1 (Biotin) (ab7815) at 0.5ug/ml

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors