abcam

Product datasheet

Biotin Anti-TARC/CCL17 antibody ab84279

Overview

Product name Biotin Anti-TARC/CCL17 antibody

DescriptionBiotin Rabbit polyclonal to TARC/CCL17

Host species Rabbit

Conjugation Biotin

Tested applications Suitable for: Sandwich ELISA, WB

Species reactivity Reacts with: Human

Immunogen Recombinant full length protein corresponding to Human TARC/CCL17.

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Lyophilized:Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a

concentration of 0.1-1.0 mg/ml.

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze /

thaw cycle.

Storage buffer Constituent: PBS

Purification notes ab84279 was purified by affinity chromatography and then biotinylated.lt is sterile filtered.

Clonality Polyclonal

Isotype IgG

Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab84279 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

1

Application	Abreviews	Notes
Sandwich ELISA		Use a concentration of 0.25 - 1 µg/ml.
WB		Use a concentration of 0.1 - 0.2 µg/ml. Predicted molecular weight: 11 kDa.

Target		
Function	Chemotactic factor for T-lymphocytes but not monocytes or granulocytes. May play a role in T-cell development in thymus and in trafficking and activation of mature T-cells. Binds to CCR4.	
Tissue specificity	Expressed at high levels in thymus and at low levels in the lung, colon and small intestine.	
Sequence similarities	Belongs to the intercrine beta (chemokine CC) family.	
Cellular localization	Secreted.	

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors