abcam

Product datasheet

Anti-CD4 antibody [OKT4] ab218628

Overview

Product name Anti-CD4 antibody [OKT4]

Description Mouse monoclonal [OKT4] to CD4

Host species Mouse

Tested applications Suitable for: Flow Cyt
Species reactivity Reacts with: Human

Immunogen The details of the immunogen for this antibody are not available.

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Storage buffer pH: 7.20

Preservative: 0.09% Sodium azide

 $Constituents: 0.87\% \ Sodium \ chloride, 0.12\% \ Monobasic \ dihydrogen \ sodium \ phosphate$

Purity Protein A purified

Purification notes ab218628 was purified from tissue culture supernatant via affinity chromatography. >90% pure as

determined by SDS-PAGE analysis

Clonality Monoclonal

Clone number OKT4

Isotype IgG2b

Light chain type kappa

A-----

1

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab218628 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use at an assay dependent concentration.

Target

Function Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell

activation. Induces the aggregation of lipid rafts.

Sequence similarities Contains 3 lg-like C2-type (immunoglobulin-like) domains.

Contains 1 lg-like V-type (immunoglobulin-like) domain.

Post-translational modifications

Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.

Cellular localization Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein

that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation.

Cell surface expression is also down-modulated by HIV-1 Envelope polyprotein gp160 that

interacts with, and sequesters CD4 in the endoplasmic reticulum.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors