


Anti-CD48 antibody [MEM-102] ab9185

[2 References](#) [2 Images](#)

Overview

Product name	Anti-CD48 antibody [MEM-102]
Description	Mouse monoclonal [MEM-102] to CD48
Host species	Mouse
Tested applications	Suitable for: Flow Cyt
Species reactivity	Reacts with: Human Predicted to work with: Non human primates 
Immunogen	Tissue, cells or virus corresponding to Human CD48. Burkitt's lymphoma cell line Raji
Positive control	Flow Cyt: Human peripheral blood.
General notes	<p>This antibody is more than 95 % pure (by PAGE).</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: PBS
Purity	Protein A purified
Purification notes	Purified from TCS. Purity >95% by SDS-PAGE.
Clonality	Monoclonal
Clone number	MEM-102
Myeloma	unknown

Isotype	IgG1
Light chain type	unknown

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab9185 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use a concentration of 2 µg/ml.


Target

Function	Ligand for CD2. Might facilitate interaction between activated lymphocytes. Probably involved in regulating T-cell activation.
Sequence similarities	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Cellular localization	Cell membrane.

Images



Flow cytometry analysis of human peripheral whole blood labeling CD48 with ab9185 at 3 µg/mL (GAM APC).



Flow cytometry analysis of human peripheral whole blood labeling CD48 positive lymphocytes with ab9185 at 3 µg/mL (GAM APC) from neutrophil granulocytes (black-dashed).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors