

## Product datasheet

### Anti-CD79α antibody [EP3618] ab79414

KO VALIDATED

Recombinant

RabMAb


[5 References](#) [16 Images](#)

#### Overview

<b>Product name</b>	Anti-CD79a antibody [EP3618]
<b>Description</b>	Rabbit monoclonal [EP3618] to CD79a
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> ICC/IF, WB, IHC-P <b>Unsuitable for:</b> IP
<b>Species reactivity</b>	<b>Reacts with:</b> Human
<b>Immunogen</b>	Synthetic peptide within Human CD79a aa 50-150 (extracellular). The exact sequence is proprietary.
<b>Positive control</b>	Raji cell, Daudi cell, human tonsil tissue and human spleen tissue lysates. Raji cells. Human tonsil tissue. WB: Wild-type Raji cell lysate and Daudi cell lysate
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb® patents</a>.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

#### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
<b>Dissociation constant (K<sub>D</sub>)</b>	K <sub>D</sub> = 1.16 x 10 <sup>-10</sup> M



[Learn more about Kp](#)

<b>Storage buffer</b>	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture supernatant
<b>Purity</b>	Tissue culture supernatant
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EP3618
<b>Isotype</b>	IgG

## Applications

**The Abpromise guarantee** Our [Abpromise guarantee](#) covers the use of ab79414 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/500.
WB		1/5000 - 1/20000. Detects a band of approximately 44-48 kDa (predicted molecular weight: 25 kDa).
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

**Application notes** Is unsuitable for IP.

## Target


<b>Function</b>	Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of immature B cells.
<b>Tissue specificity</b>	B-cells.
<b>Involvement in disease</b>	Defects in CD79A are the cause of agammaglobulinemia type 3 (AGM3) [MIM:613501]. It is a primary immunodeficiency characterized by profoundly low or absent serum antibodies and low or absent circulating B cells due to an early block of B-cell development. Affected individuals develop severe infections in the first years of life. Note=Two different mutations, one at the splice donor site of intron 2 and the other at the splice acceptor site for exon 3, have been identified. Both mutations give rise to a truncated protein.
<b>Sequence similarities</b>	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 ITAM domain.
<b>Post-translational modifications</b>	Phosphorylated on tyrosine, serine and threonine residues upon B-cell activation. Phosphorylation of tyrosine residues by Src-family kinases is an early and essential feature of the BCR signaling

cascade. The phosphorylated tyrosines serve as docking sites for SH2-domain containing kinases, leading to their activation which in turn leads to phosphorylation of downstream targets. Phosphorylation of serine and threonine residues may prevent subsequent tyrosine phosphorylation.

## Cellular localization

Cell membrane. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts.

## Images



**All lanes :** Anti-CD79a antibody [EP3618] (ab79414) at 1/5000 dilution

**Lane 1 :** Wild-type Raji cell lysate

**Lane 2 :** CD79A knockout Raji cell lysate

**Lane 3 :** Daudi cell lysate

**Lane 4 :** HAP1 cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 25 kDa

**Observed band size:** 40-50 kDa


False colour image of Western blot: Anti-CD79a antibody [EP3618] staining at 1/5000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab79414 was shown to bind specifically to CD79a. A band was observed at 40-50 kDa in wild-type Raji cell lysates with no signal observed at this size in CD79A knockout cell line [ab274911](#) (knockout cell lysate [ab281361](#)). To generate this image, wild-type and CD79A knockout Raji cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ([ab216776](#)) at 1/20000

dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody [EP3618] (ab79414)

ab79414 at 1/100 dilution staining CD79a in paraffin-embedded human tonsil tissue by Immunohistochemistry. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-CD79a antibody [EP3618] (ab79414)

**All lanes :** Anti-CD79a antibody [EP3618] (ab79414) at 1/10000 dilution

**Lane 1 :** Raji cell lysate

**Lane 2 :** Daudi cell lysate

**Lane 3 :** Human tonsil tissue lysate

**Lane 4 :** Human spleen tissue lysate


Lysates/proteins at 10 µg per lane.

#### Secondary

**All lanes :** HRP labelled goat anti-rabbit at 1/2000 dilution

**Predicted band size:** 25 kDa


**Observed band size:** 44-48 kDa



Immunocytochemistry/ Immunofluorescence - Anti-CD79a antibody [EP3618] (ab79414)


Immunocytochemistry/Immunofluorescence analysis of RAMOS (human Burkitt's lymphoma) cells labelling CD79a with purified ab79414 at 1/500. Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. Nuclei were counterstained with DAPI (blue).

Secondary Only Control: PBS was used instead of the primary antibody as the negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody [EP3618] (ab79414)

ab79414 showing positive staining in Normal spleen tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody [EP3618] (ab79414)

ab79414 showing positive staining in Normal tonsil tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody  
[EP3618] (ab79414)

ab79414 showing negative staining in Normal brain tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody  
[EP3618] (ab79414)


ab79414 showing negative staining in Normal colon tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody  
[EP3618] (ab79414)


ab79414 showing negative staining in Normal heart tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody  
[EP3618] (ab79414)

ab79414 showing negative staining in Normal brain tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody  
[EP3618] (ab79414)


ab79414 showing positive staining in Normal tonsil tissue. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody  
[EP3618] (ab79414)

ab79414 showing negative staining in Normal colon tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody [EP3618] (ab79414)

ab79414 showing negative staining in Normal heart tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD79a antibody [EP3618] (ab79414)

ab79414 showing positive staining in Normal spleen tissue.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



OL-RD Scanning - Anti-CD79a antibody [EP3618] (ab79414)

Equilibrium disassociation constant ( $K_D$ )

Learn more about  $K_D$

[Click here to learn more about  \$K\_D\$](#)



### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-CD79a antibody [EP3618] (ab79414)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors