

Anti-Cytomegalovirus pp65 antibody [CH12] ab53489

8 References

Overview

Product name	Anti-Cytomegalovirus pp65 antibody [CH12]
Description	Mouse monoclonal [CH12] to Cytomegalovirus pp65
Host species	Mouse
Specificity	ab53489 is reactive with pp65 (UL83) of Cytomegalovirus.
Tested applications	Suitable for: ICC/IF, WB
Species reactivity	Reacts with: Human cytomegalovirus
Immunogen	Tissue, cells or virus corresponding to Cytomegalovirus pp65. HCMV AD169 IC Extract
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7.40 Constituent: PBS
Purity	Protein G purified
Clonality	Monoclonal
Clone number	CH12
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab53489 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use a concentration of 10 µg/ml.
WB		Use a concentration of 10 µg/ml.

Target

Relevance

Cytomegalovirus is a member of the herpes virus group, which includes herpes simplex virus types 1 and 2, varicella zoster virus (which causes chicken pox), and Epstein Barr virus (which causes infectious mononucleosis). These viruses share a characteristic ability to remain dormant within the body over a long period. CMV viral genes are co-ordinately expressed in groups at various times after infection. Early viral proteins are expressed in the nucleus of infected cells within 3 to 24 hours of infection prior to the commencement of viral DNA replication. This is followed by expression of the early intermediate genes, which encode enzymes required for viral DNA replication. After 48 to 72 hours, a number of late viral antigens may be demonstrated in the nuclei and cytoplasm of infected cells. pp65 is a 65kD phosphorylated glycoprotein and is the most abundant of the late antigens.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors