

Anti-DPP9 antibody - Catalytic domain ab42080

★★★★★ [7 Abreviews](#) [26 References](#) [4 Images](#)

Overview

Product name	Anti-DPP9 antibody - Catalytic domain
Description	Rabbit polyclonal to DPP9 - Catalytic domain
Host species	Rabbit
Specificity	This antibody recognises all three forms of DPP9. This antibody does not recognise other DPP family members.
Tested applications	Suitable for: ICC/IF, WB, IHC-P
Species reactivity	Reacts with: Cow, Human
Immunogen	Synthetic peptide corresponding to Human DPP9. (Peptide available as ab44599)
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.05% Sodium azide Constituents: PBS, 50% Glycerol, 2.9% Sodium chloride
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab42080 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF	★★★★★ (1)	Use a concentration of 1 µg/ml.
WB	★★★★★ (3)	1/1000 - 1/5000. Predicted molecular weight: 98 kDa. Recommended dilution 1/1000 with colourimetric substrates, 1/5000 dilution with chemiluminescent substrates. Dilution optimised using Chromogenic detection.
IHC-P	★★★★★ (3)	1/50.

Target

Function	Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2.
Tissue specificity	Ubiquitously expressed, with highest levels in liver, heart and muscle, and lowest levels in brain.
Sequence similarities	Belongs to the peptidase S9B family. DPPIV subfamily.
Cellular localization	Cytoplasm > cytosol.

Images




All lanes : Anti-DPP9 antibody - Catalytic domain (ab42080) at 1/1000 dilution

Lane 1 : Cell media from human prostate cancer DU145 cells (treated with SF)

Lane 2 : Cell media from human prostate cancer DU145 cells (treated with RA)

Lane 3 : Cell media from human prostate cancer DU145 cells (treated with IL1-beta)

Predicted band size: 98 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DPP9 antibody - Catalytic domain (ab42080)

Ab42080 staining Human normal placenta. Staining is localised to the membrane and cytoplasm.

Left panel: with primary antibody at 4 ug/ml. Right panel: isotype control.


Sections were stained using an automated system DAKO Autostainer Plus, at room temperature. Sections were rehydrated and antigen retrieved with the Dako 3-in-1 AR buffers EDTA pH 9.0 in a DAKO PT Link. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 minutes. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 minutes and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DPP9 antibody - Catalytic domain (ab42080)

This image is a courtesy of an Abreview submitted by Veronique Dubois

ab42080 staining DPP9 in bovine testicular tissue section by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue underwent paraformaldehyde fixation before heat mediated antigen retrieval in Tris/EDTA buffer pH 9.0 and then the sample was blocked with 3% hydrogen peroxide in TBS for 10 minutes at 25°C. The sections were treated for 10 min with 3% H₂O₂ in TBS to block endogenous peroxidase activity and incubated subsequently for 30 min with 10% nonimmune goat serum to minimize nonspecific antibody binding. The primary antibody (1/50) was incubated with sample in TBS, 0.3% Triton X-100, 0.1 % BSA for 16 hours at 4°C. A Biotin conjugated goat polyclonal to rabbit IgG was used at dilution 1/200 as secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-DPP9 antibody - Catalytic domain (ab42080)

ICC/IF image of ab42080 stained Hek293 cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab42080, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors