

Product datasheet

Anti-ds DNA antibody [35I9 DNA] - BSA and Azide free ab27156

★★★★★ [10 Abreviews](#) [99 References](#) [1 Image](#)

Overview

Product name	Anti-ds DNA antibody [35I9 DNA] - BSA and Azide free
Description	Mouse monoclonal [35I9 DNA] to ds DNA - BSA and Azide free
Host species	Mouse
Specificity	Primarily Double stranded DNA. Measurements by immuno-CE yielded K_D 's of 0.71 μ M and 0.09 μ M, for the interaction of this antibody with ss- and dsDNA, respectively. Strong reactivity with both ss- and dsDNA has been observed on dotblots as well as very weak reactivity with RNA.
Tested applications	Suitable for: IHC-P
Species reactivity	Reacts with: Human
Immunogen	The details of the immunogen for this antibody are not available.
Positive control	IHC-P: human skin tissue sections.
General notes	<p>This antibody has been shown to be useful, in the detection of dsDNA in fx. <i>Crithidia luciliae</i> -a monoflagellate protozoan,containing a giant mitochondrion. Very good reactivity both with dsDNA and ssDNA on NC-dotblots has been observed. The minimal size for DNA binding for this antibody is >16 bases and there is a inverse proportionality between binding and ionic strength. Abcam recommended secondaries - Goat Anti-Mouse HRP (ab205719) and Goat Anti-Mouse Alexa Fluor® 488 (ab150113). Or search our wide range of secondary antibodies for use with your experiment.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Storage buffer	pH: 7.20 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Primary antibody notes	This antibody has been shown to be useful, using immunofluorescence, in the detection of dsDNA in fx. <i>Crithidia luciliae</i> -a monoflagellate protozoan, containing a giant mitochondrion. The antibody can be used for easy quantification of in vitro cell proliferation by ELISA and has successfully been used for monitoring cytotoxicity and apoptosis as well as ELISA-based co-culture angiogenesis and proliferation assays. Very good reactivity both with dsDNA and ssDNA on NC-dotblots has been observed. The minimal size for DNA binding for this antibody is >16 bases and there is a inverse proportionality between binding and ionic strength.
Clonality	Monoclonal
Clone number	3519 DNA
Myeloma	x63-Ag8.653
Isotype	IgG2a
Light chain type	kappa

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab27156 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★★★★★ (1)	Use a concentration of 0.05 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Target

Relevance	dsDNA (double stranded deoxyribonucleic acid) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3',5'-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.
Cellular localization	Nuclear

Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ds DNA antibody [3519 DNA] - BSA and Azide free (ab27156)

IHC image of ds DNA staining in a section of formalin-fixed paraffin-embedded normal human skin* performed on a Leica BOND™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab27156, 0.05ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors