

Product datasheet

Anti-EpCAM antibody [EPR677(2)] α b124825

Recombinant RabMAb

★☆☆☆☆ **1 Abreviews** **10 References** [13 Images](#)

Overview

Product name	Anti-EpCAM antibody [EPR677(2)]
Description	Rabbit monoclonal [EPR677(2)] to EpCAM
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P Unsuitable for: Flow Cyt, ICC/IF or IP
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide within Human EpCAM aa 50-150. The exact sequence is proprietary. Database link: P16422
Positive control	WB: HCT116, A431 and HT29 cell lysates. IHC-P: Human colon adenocarcinoma, colon, endometrial adenocarcinoma and stomach adenocarcinoma tissues.
General notes	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	pH: 7.2 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
Purity	Protein A purified
Clonality	Monoclonal

Clone number EPR677(2)

Isotype IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab124825 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000 - 1/10000. Detects a band of approximately 39 kDa (predicted molecular weight: 35 kDa). Western blots of mouse and rat samples are negative.
IHC-P	★☆☆☆☆ (1)	1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. Heat up to 98 degrees C, below boiling, and then let cool for 10-20 min.

Application notes Is unsuitable for Flow Cyt, ICC/IF or IP.

Target

Function May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

Tissue specificity Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC). Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Involvement in disease Defects in EPCAM are the cause of diarrhea type 5 (DIAR5) [MIM:613217]. It is an intractable diarrhea of infancy characterized by villous atrophy and absence of inflammation, with intestinal epithelial cell dysplasia manifesting as focal epithelial tufts in the duodenum and jejunum. Defects in EPCAM are a cause of hereditary non-polyposis colorectal cancer type 8 (HNPCC8) [MIM:613244]. HNPCC is a disease associated with marked increase in cancer susceptibility. It is characterized by a familial predisposition to early-onset colorectal carcinoma (CRC) and extra-colonic tumors of the gastrointestinal, urological and female reproductive tracts. HNPCC is reported to be the most common form of inherited colorectal cancer in the Western world. Clinically, HNPCC is often divided into two subgroups. Type I is characterized by hereditary predisposition to colorectal cancer, a young age of onset, and carcinoma observed in the proximal colon. Type II is characterized by increased risk for cancers in certain tissues such as the uterus, ovary, breast, stomach, small intestine, skin, and larynx in addition to the colon. Diagnosis of classical HNPCC is based on the Amsterdam criteria: 3 or more relatives affected by colorectal cancer, one a first degree relative of the other two; 2 or more generation affected; 1 or more colorectal cancers presenting before 50 years of age; exclusion of hereditary polyposis syndromes. The term 'suspected HNPCC' or 'incomplete HNPCC' can be used to describe families who do not or only partially fulfill the Amsterdam criteria, but in whom a genetic basis for colon cancer is strongly suspected. Note=HNPCC8 results from heterozygous deletion of 3-prime

exons of EPCAM and intergenic regions directly upstream of MSH2, resulting in transcriptional read-through and epigenetic silencing of MSH2 in tissues expressing EPCAM.

Sequence similarities

Belongs to the EPCAM family.

Contains 1 thyroglobulin type-1 domain.

Post-translational modifications


Hyperglycosylated in carcinoma tissue as compared with autologous normal epithelia.

Glycosylation at Asn-198 is crucial for protein stability.

Cellular localization

Lateral cell membrane. Cell junction > tight junction. Co-localizes with CLDN7 at the lateral cell membrane and tight junction.


Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825 showing positive staining in Normal kidney tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825, at 1/100 dilution, staining EpCAM in paraffin-embedded Human colon adenocarcinoma tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-EpCAM antibody [EPR677(2)]
(ab124825)

All lanes : Anti-EpCAM antibody [EPR677(2)] (ab124825) at 1/1000 dilution

Lane 1 : HCT116 cell lysate

Lane 2 : A431 cell lysate


Lane 3 : HT29 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat anti-Rabbit HRP at 1/2000 dilution


Predicted band size: 35 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825, at 1/100 dilution, staining EpCAM in paraffin-embedded Human colon tissue by Immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825, at 1/100 dilution, staining EpCAM in paraffin-embedded Human endometrial adenocarcinoma tissue by Immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825 showing positive staining in Urinary bladder transitional carcinoma tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825 showing negative staining in Normal brain tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825 showing negative staining in Glioma tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825 showing negative staining in Skeletal muscle tissue.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825 showing negative staining in Normal tonsil tissue.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

ab124825, at 1/100 dilution, staining EpCAM in paraffin-embedded Human stomach adenocarcinoma tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Tissue Microarray (TMA) data for ab124825					
Normal tissue samples			Malignant tissue samples		
Human cardiac muscle	x	Human placenta	x	Human glioma	x
Human cerebrum	x	Human skeletal muscle	x	Human hepatocellular carcinoma	x
Human colon	✓	Human skin	✓	Human lung carcinoma	✓
Human endometrium	✓	Human spleen	x	Human ovarian carcinoma	✓
Human kidney	✓	Human stomach	x	Human pancreatic carcinoma	x
Human liver	x	Human testis	x	Human prostatic hyperplasia	x
Human lung	x	Human thyroid	x	Human thyroid carcinoma	x
Human mammary gland	✓	Human tonsil	x		
Human pancreas	✓				

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EpCAM antibody [EPR677(2)] (ab124825)

Tissue Microarrays stained for "Anti-EpCAM antibody [EPR677(2)]" using "ab124825" in immunohistochemical analysis. This table provides a detailed overview of positive (tick mark) and negative (cross mark) staining per sample type tested. The sections were pre-treated using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) for 20 minutes. The sections were incubated with ab124825 for 30 mins at room temperature followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-EpCAM antibody [EPR677(2)] (ab124825)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors