

Product datasheet

Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free ab194979

KO VALIDATED Recombinant RabMAB

[7 References](#) [12 Images](#)

Overview


Product name	Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free
Description	Rabbit monoclonal [EP1045Y] to ErbB2 / HER2 - BSA and Azide free
Host species	Rabbit
Specificity	ab134182 detects ErbB 2 phosphorylated at Tyr1248 as well as unphosphorylated ErbB 2. Mouse species is recommended based on WB results, we do not guarantee IHC-P for mouse.
Tested applications	Suitable for: WB, ICC/IF, IHC-P, IP Unsuitable for: Flow Cyt
Species reactivity	Reacts with: Mouse, Human
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: HeLa, SKBR-3, Wild-type HCT 116 and Wild-type A549 cell lysates; IHC-P: Human breast carcinoma tissue; ICC/IF: SKBR cells; IP: HeLa.
General notes	<p>ab194979 is the carrier-free version of ab134182.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Dissociation constant (K _D)	K _D = 3.00 x 10 ⁻¹¹ M



Storage buffer	pH: 7.2 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EP1045Y
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab194979 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Detects a band of approximately 185 kDa (predicted molecular weight: 137 kDa). Please check the parent abID, ab134182 , for more information on dilutions.
ICC/IF		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
IP		Use at an assay dependent concentration.

Application notes Is unsuitable for Flow Cyt.

Target

Function	<p>Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC. Involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.</p>
Tissue specificity	<p>Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.</p>
Involvement in disease	<p>Hereditary diffuse gastric cancer Glioma Ovarian cancer Lung cancer Gastric cancer</p> <p>Chromosomal aberrations involving ERBB2 may be a cause gastric cancer. Deletions within 17q12 region producing fusion transcripts with CDK12, leading to CDK12-ERBB2 fusion leading to truncated CDK12 protein not in-frame with ERBB2.</p>
Sequence similarities	<p>Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. Contains 1 protein kinase domain.</p>
Post-translational modifications	<p>Autophosphorylated. Autophosphorylation occurs in trans, i.e. one subunit of the dimeric receptor phosphorylates tyrosine residues on the other subunit (Probable). Ligand-binding increases phosphorylation on tyrosine residues (PubMed:27134172). Signaling via SEMA4C promotes phosphorylation at Tyr-1248 (PubMed:17554007). Dephosphorylated by PTPN12 (PubMed:27134172).</p>
Cellular localization	<p>Cytoplasm. Nucleus and Cell membrane. Cytoplasm, perinuclear region. Nucleus. Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1.</p>

Images



Western blot - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

All lanes : Anti-ErbB2 / HER2 antibody [EP1045Y] ([ab134182](#)) at 1/500 dilution

Lane 1 : Wild-type MCF7 cell lysate at 32 µg

Lane 2 : ERBB2 knockout MCF7 cell lysate at 32 µg

Lane 3 : A549 cell lysate at 16 µg


Performed under reducing conditions.

Predicted band size: 137 kDa

Observed band size: 180 kDa

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab134182](#)).

Western blot: Anti-ErbB2 / HER2 antibody [EP1045Y] staining at 1/500 dilution, shown in black; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab134182](#) was shown to bind specifically to ErbB2 / HER2. A band was observed at 180 kDa in wild-type MCF7 cell lysates with no signal observed at this size in ERBB2 knockout cell line [ab286260](#) (knockout cell lysate AB300208). To generate this image, wild-type and ERBB2 knockout MCF7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature and washed again four times. Secondary antibodies used were HRP conjugated Goat anti-Rabbit (H+L) and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution. This blot was developed with an ultra high-sensitivity ECL substrate kit and imaged with 20 minutes exposure time.



All lanes : Anti-ErbB2 / HER2 antibody [EP1045Y] ([ab134182](#)) at 1/1000 dilution

Lane 1 : Wild-type A549 cell lysate

Lane 2 : ERBB2 knockout A549 cell lysate

Lane 3 : HeLa cell lysate

Lane 4 : SK-BR-3 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.


Predicted band size: 137 kDa

Observed band size: 180 kDa

False colour image of Western blot: Anti-ErbB2 / HER2 antibody [EP1045Y] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab134182](#) was shown to bind specifically to ErbB2 / HER2. A band was observed at 180 kDa in wild-type A549 cell lysates with no signal observed at this size in ERBB2 knockout cell line. To generate this image, wild-type and ERBB2 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG

H&L 680RD at 1/20000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab134182](#)).



All lanes : Anti-ErbB2 / HER2 antibody [EP1045Y] ([ab134182](#)) at 1/1000 dilution

Lane 1 : Wild-type HCT 116 cell lysate

Lane 2 : ERBB2 knockout HCT 116 cell lysate

Lane 3 : HeLa cell lysate

Lane 4 : SK-BR-3 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.


Predicted band size: 137 kDa

Observed band size: 180 kDa

False colour image of Western blot: Anti-ErbB2 / HER2 antibody [EP1045Y] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab134182](#) was shown to bind specifically to ErbB2 / HER2. A band was observed at 180 kDa in wild-type HCT 116 cell lysates with no signal observed at this size in ERBB2 knockout cell line. To generate this image, wild-type and ERBB2 knockout HCT 116 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in fluorescent western blot (TBS-based) blocking solution before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and

sodium azide ([ab134182](#)).



Western blot - Anti-ErbB2 / HER2 antibody
[EP1045Y] - BSA and Azide free (ab194979)

All lanes : Anti-ErbB2 / HER2 antibody [EP1045Y] ([ab134182](#)) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate


Lane 2 : ERBB2 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 137 kDa


False colour image of Western blot: Anti-ErbB2 / HER2 antibody [EP1045Y] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab134182](#) was shown to bind specifically to ErbB2 / HER2. A band was observed at 180 kDa in wild-type HeLa cell lysates with no signal observed at this size in ERBB2 knockout cell line [ab255387](#) (knockout cell lysate [ab263758](#)). To generate this image, wild-type and ERBB2 knockout HeLa cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast carcinoma tissue sections labeling ErbB2 / HER2 with Purified **ab134182** at 1:1600 dilution (0.68 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab134182**).



Western blot - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979) + SKBR-3 (human mammary gland adenocarcinoma) whole cell lysate at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**)


Predicted band size: 137 kDa

Observed band size: 185 kDa

Exposure time: 3 seconds

Blocking buffer and concentration: 5% NFDm/TBST

Diluting buffer and concentration: 5% NFDm/TBST



Immunoprecipitation - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

ab134182 (purified) at 1:30 dilution (2µg) immunoprecipitating ErbB2 / HER2 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg


Lane 2 (+): **ab134182** & HeLa whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab134182** in HeLa whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab134182**).




Immunocytochemistry/ Immunofluorescence - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Immunocytochemistry/Immunofluorescence analysis of SK-BR-3 (human mammary gland adenocarcinoma) labelling ErbB2 / HER2 with purified **ab134182** at 1/125. Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100. An Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody (Ab150077). Nuclei counterstained with DAPI (blue).

Control: PBS only

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab134182**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue labelling ErbB2 / HER2 with **ab134182** at 1/100 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab134182**).


Heat mediated antigen retrieval was performed with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Immunofluorescent analysis of SKBR cells labelling ErbB2 / HER2 with **ab134182** at 1/250 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab134182**).



SPR Scanning - Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab134182**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-ErbB2 / HER2 antibody [EP1045Y] - BSA and Azide free (ab194979)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

-
- Replacement or refund for products not performing as stated on the datasheet
 - Valid for 12 months from date of delivery
 - Response to your inquiry within 24 hours

 - We provide support in Chinese, English, French, German, Japanese and Spanish
 - Extensive multi-media technical resources to help you
 - We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors