abcam

Product datasheet

FITC Anti-IgE antibody [BE5] ab49434

1 References

Overview

Product name FITC Anti-lgE antibody [BE5]

Description FITC Mouse monoclonal [BE5] to IgE

Host species Mouse

Conjugation FITC. Ex: 493nm, Em: 528nm

Tested applications Suitable for: Flow Cyt
Species reactivity Reacts with: Human

Immunogen Full length native protein (purified) corresponding to Human IgE.

General notes The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum

conditions. The reagent is free of unconjugated FITC.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C.

Storage buffer pH: 7.4

Preservative: 0.097% Sodium azide Constituents: PBS, 0.2% BSA

Purity Size exclusion

Clonality Monoclonal

Clone number BE5 Isotype IgG1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab49434 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use at an assay dependent concentration. ab91356 - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.

Target

Relevance

IgE is the class of antibodies produced in the lungs, skin, and mucous membranes. It may protect against parasite invasion, but it is a major factor in allergic reactions. The antigen-specific IgE interacts with mast cells and eosinophils, triggers the release of histamine, leukotrienes and other substances that lead to the itching, sneezing and congestion of allergies - and the life threatening respiratory distress of asthma and anaphylactic shock.

Cellular localization

Cell Membrane

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors