

Product datasheet

Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free ab197544


KO VALIDATED

Recombinant

RabMAb

15 Images

Overview

Product name	Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free
Description	Rabbit monoclonal [EP2775Y] to Galectin 3 - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: IHC-P, WB, Flow Cyt (Intra), ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Human Predicted to work with: Pig 
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: A375, HeLa, SW480, A431, RAW264.7 and NIH/3T3, C6, Wild-type A549 and MCF7 cell lysates. ICC/IF: Panc-1 and HT-29 cells, RAW264.7 cells and C6 cells . IHC-P: Human lung squamous carcinoma and human thyroid carcinoma tissues. Flow cyt (intra): RAW264.7 cells and C6 cells.
General notes	<p>ab197544 is the carrier-free version of ab76245.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	pH: 7.20 Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EP2775Y
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab197544 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Predicted molecular weight: 26 kDa.
Flow Cyt (Intra)		Use at an assay dependent concentration. <u>ab199376</u> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.

Target

Function	Galactose-specific lectin which binds IgE. May mediate with the alpha-3, beta-1 integrin the stimulation by CSPG4 of endothelial cells migration. Together with DMBT1, required for terminal differentiation of columnar epithelial cells during early embryogenesis.
Tissue specificity	A major expression is found in the colonic epithelium. It is also abundant in the activated macrophages.
Sequence similarities	Contains 1 galectin domain.
Cellular localization	Nucleus. Cytoplasmic in adenomas and carcinomas. May be secreted by a non-classical secretory pathway and associate with the cell surface.

Images



All lanes : Anti-Galectin 3 antibody [EP2775Y] ([ab76245](#)) at 1/5000 dilution

Lane 1 : Wild-type A549 cell lysate

Lane 2 : LGALS3 knockout A549 cell lysate

Lane 3 : MCF7 cell lysate

Lane 4 : LNCaP cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680R at 1/20000 dilution


Performed under reducing conditions.

Predicted band size: 26 kDa

Observed band size: 30 kDa

False colour image of Western blot: Anti-Galectin 3 antibody [EP2775Y] staining at 1/5000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab76245](#) was shown to bind specifically to Galectin 3. A band was observed at 30 kDa in wild-type A549 cell lysates with no signal observed at this size in LGALS3 knockout cell line. To generate this image, wild-type and LGALS3 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76245](#)).



Flow Cytometry (Intracellular) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76245](#)).


Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized C6 (rat glial tumor glial cell) cells labelling Galectin 3 with [ab76245](#) at 1:50 dilution (1µg)/ Red compared with a Rabbit monoclonal IgG ([ab172730](#)) / Black isotype control and an unlabelled control (Cell without incubation with primary antibody and secondary antibody (Blue)). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, [ab150081](#)) at 1/2000 dilution was used as the secondary antibody.



Flow Cytometry (Intracellular) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76245](#)).


Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) cells labelling Galectin 3 with [ab76245](#) at 1:50 dilution (1µg)/ Red compared with a Rabbit monoclonal IgG ([ab172730](#)) / Black isotype control and an unlabelled control (Cell without incubation with primary antibody and secondary antibody (Blue)). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, [ab150081](#)) at 1/2000 dilution was used as the secondary antibody.



This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized C6 cells labeling Galectin 3 with **ab76245** at 1/100 dilution, followed by **ab150081** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed secondary antibody at 1/1000 dilution. **ab195889** Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution. The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is **ab150081** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution.


Confocal image showing cytoplasmic and weak nuclear staining in C6 cell line.



This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized RAW 264.7 cells labeling Galectin 3 with **ab76245** at 1/100 dilution, followed by **ab150081** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed secondary antibody at 1/1000 dilution. **ab195889** Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution. The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is **ab150081** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution.

Confocal image showing nuclear and cytoplasmic staining in RAW 264.7 cell line.



All lanes : Anti-Galectin 3 antibody [EP2775Y] (**ab76245**) at 1/1000 dilution

Lane 1 : RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate

Lane 2 : Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate

Lane 3 : A431 (human epidermoid carcinoma epithelial cell) whole cell lysate

Lane 4 : HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 5 : SK-N-MC (human brain epithelial cell) whole cell lysate

Lane 6 : LNCaP (human prostate carcinoma epithelial cell) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Predicted band size: 26 kDa


Observed band size: 26 kDa

Exposure time: 26 seconds

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).

Blocking and diluting buffer and concentration: 5% NFDm/TBST,

ab129002 was used as a loading control for Vinculin.



Western blot - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

All lanes : Anti-Galectin 3 antibody [EP2775Y] ([ab76245](#)) at 1/1000 dilution

Lane 1 : C6 (rat glial tumor glial cell) whole cell lysate

Lane 2 : RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate

Lane 3 : NIH/3T3 (mouse embryonic fibroblast) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 26 kDa

Observed band size: 26 kDa

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab76245](#)).

Blocking and diluting buffer and concentration: 5% NFDm/TBST.

Exposure time: Lane 1: 180 seconds, Lanes 2 and 3: 15 seconds.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

This image is courtesy of an Abreview submitted by Carl Hobbs.

Immunohistochemical analysis of formaldehyde fixed mouse lung tissue sections labelling Galectin 3 with **ab76245** at a dilution of 1/6000. The secondary antibody used was biotin conjugated goat anti rabbit IgG at a dilution of 1/300. Antigen retrieval was heat mediated using citric acid.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).



Flow Cytometry (Intracellular) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labeling Galectin 3 with purified **ab76245** at 1/50 dilution (10ug/mL) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).




Immunocytochemistry/ Immunofluorescence - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

Immunocytochemistry/Immunofluorescence analysis of HT-29 cells labelling Galectin with purified **ab76245** at 1/1000. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/500) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/100) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).



Immunocytochemistry/ Immunofluorescence - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

ICC/IF image of unpurified **ab76245** stained Panc-1 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody **ab76245** at 10µg/ml overnight at +4°C. The secondary antibody (pseudo-colored green) was Alexa Fluor® 488 goat anti-rabbit (**ab150081**) IgG (H+L) preadsorbed, used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (pseudo-colored red) at a 1/200 dilution for 1h at room temperature. DAPI was used to stain the cell nuclei (pseudo-colored blue) at a concentration of 1.43µM for 1 hour at room temperature.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).



Flow Cytometry (Intracellular) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

Overlay histogram showing THP1 cells stained with unpurified **ab76245** (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% human serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab76245**, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung squamous carcinoma tissue labelling Galectin 3 with unpurified **ab76245**.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab76245**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

This IHC data was generated using the same anti-Galectin 3 antibody clone [EP2775Y] in a different buffer formulation (cat# **ab76245**).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human thyroid carcinoma tissue labelling Galectin 3 with purified **ab76245** at 1/250. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Why choose a recombinant antibody?



Anti-Galectin 3 antibody [EP2775Y] - BSA and Azide free (ab197544)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors