

Product datasheet

Anti-HIV1 gp41 antibody ab20890

Overview

Product name	Anti-HIV1 gp41 antibody
Description	Goat polyclonal to HIV1 gp41
Host species	Goat
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Human immunodeficiency virus
Immunogen	Recombinant (P. Pastoris) ectodomain of gp41 (glycosylated)
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.1% Sodium azide Constituent: 0.0268% PBS
Purity	Ion Exchange Chromatography
Purification notes	Ab20890 is purified by sodium sulfate precipitation and ion-exchange chromatography. The antiserum has not been adsorbed to remove anti-beta-galactosidase activity.
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab20890 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/100. Predicted molecular weight: 41 kDa. Major band seen in WB at 41kDa and a minor band at 20kDa. Dilution optimised using Chromogenic detection.

Target

Relevance

gp41/120 is the major HIV protein associated with the HIV envelope. It functions as the viral antireceptor or attachment protein. gp41 (or TM) traverses the envelope, whereas gp120 is present on the outer surface and is noncovalently attached to gp41. The precursor of gp120/41 (gp160) is synthesized in the endoplasmic reticulum and is transported via the golgi body to the cell surface. Upon activation of the envelope glycoprotein (gp120/41) by cellular receptors, gp41 undergoes conformational changes that mediate fusion of the viral and cellular membranes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors