

HRP Anti-Staphylococcus aureus antibody ab68952

Overview

Product name	HRP Anti-Staphylococcus aureus antibody
Description	HRP Rabbit polyclonal to Staphylococcus aureus
Host species	Rabbit
Conjugation	HRP
Tested applications	Suitable for: WB, ELISA
Species reactivity	Reacts with: Staphylococcus aureus
Immunogen	Tissue, cells or virus corresponding to Staphylococcus aureus. Soluble and structural antigens of Staphylococcus aureus whole bacterium; ATCC #27660.

General notes

This antibody reacts with Staphylococcus aureus in bacterial and infected tissue samples.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.002% Thimerosal (merthiolate) Constituents: 1% BSA, PBS
Purity	Proprietary Purification
Primary antibody notes	This antibody reacts with Staphylococcus aureus in bacterial and infected tissue samples.
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab68952 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/20 - 1/200.
ELISA		1/200 - 1/1000.

Target

Relevance

S.aureus binds IgG molecules by their Fc region. In serum, the bacteria will bind IgG molecules in the wrong orientation on their surface, which hinders opsonization and phagocytosis. Mutants of S. aureus lacking protein A are more efficiently phagocytosed in vitro, and mutants in infection models have diminished virulence. Due to its affinity for the Fc region of many mammalian immunoglobulins, protein A is considered a universal reagent in biochemistry and immunology.

Cellular localization

Secreted; cell wall.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors