abcam

Product datasheet

Anti-INCENP antibody [58-217] ab23956

***** 3 Abreviews 5 References 3 Images

Overview

Product name	Anti-INCENP antibody [58-217]	
Description	Mouse monoclonal [58-217] to INCENP	
Host species	Mouse	
Tested applications	Suitable for: ICC/IF, IHC-P	
Species reactivity	Reacts with: Human Does not react with: Mouse, Rat	
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.	
Positive control	IHC-P: Human anal cancer tissue	
General notes	This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com .	
	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.	
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As	

Properties

•	
Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or - 80°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
Purity	lgG fraction
Clonality	Monoclonal
Clone number	58-217
lsotype	lgG1

Applications


The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab23956 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
ICC/IF	\star \star \star \star \star (1)	Use a concentration of 1 µg/ml.
IHC-P	\star \star \star \star \star (1)	Use at an assay dependent concentration.

Target	
Function	Component of the chromosomal passenger complex (CPC), a complex that acts as a key regulator of mitosis. The CPC complex has essential functions at the centromere in ensuring correct chromosome alignment and segregation and is required for chromatin-induced microtubule stabilization and spindle assembly. Probably acts through association with AURKB or AURKC. Seems to bind directly to microtubules. Controls the kinetochore localization of BUB1.
Sequence similarities	Belongs to the INCENP family.
Cellular localization	Chromosome > centromere. Cytoplasm > cytoskeleton > spindle. Nucleus. Chromosome > centromere > kinetochore. Localizes to inner kinetochore. Localizes on chromosome arms and inner centromeres from prophase through metaphase and then transferring to the spindle midzone and midbody from anaphase through cytokinesis. Colocalizes with AURKB at mitotic chromosomes.

Images




Immunocytochemistry/ Immunofluorescence - Anti-INCENP antibody [58-217] (ab23956) ICC/IF image of ab23956 stained human HeLa cells. The cells were methanol fixed (5 min) and incubated with the antibody (ab23956, 1µg/ml) for 1h at room temperature. The secondary antibody (green) was Alexa Fluor® 488 goat anti-mouse lgG (H+L) used at a 1/1000 dilution for 1h. Image-iTTMFX Signal Enhancer was used as the primary blocking agent, 5% BSA (in TBS-T) was used for all other blocking steps. DAPI was used to stain the cell nuclei (blue). Alexa Fluor® 594 WGA was used to label plasma membranes (red).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-INCENP antibody [58-217] (ab23956)

This image is courtesy of an Abreview submitted by Antibody Solutions Ltd.



Immunocytochemistry/ Immunofluorescence - Anti-

INCENP antibody [58-217] (ab23956)

Image kindly submitted by Serena Orlando, Giulia Guarguaglini and Patrizia Lavia, from the University La Sapienza' CNR, Italy ab23956 staining human anal cancer tissue sections by IHC-P. Sections were formaldehyde fixed and subjected to heat mediated antigen retreival in citrate buffer (pH 6) prior to blocking in a commercially available blocking agent for 20 minutes at 20°C. The primary anibody was diluted 1/200 and incubated with the sample for 40 minutes at 20°C. A HRP-conjugated goat anti-mouse antibody was used as the secondary.

ab23956 staining INCENP in HeLa cells, by immunofluorescence.

optimal antibody dilution: 1mg/ml

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors