

Product datasheet

Anti-MGMT antibody [MT3.1] ab39253

KO **VALIDATED** Recombinant

★★★★☆ **6 Abreviews** **19 References** **6 Images**

Overview

Product name	Anti-MGMT antibody [MT3.1]
Description	Mouse monoclonal [MT3.1] to MGMT
Host species	Mouse
Tested applications	Suitable for: WB, IHC-P
Species reactivity	Reacts with: Human
Immunogen	Recombinant full length protein corresponding to Human MGMT.
Positive control	IHC-P: Human tonsil, prostate carcinoma and lung carcinoma tissue. WB: MCF7, HeLa, Jurkat and HCT116 whole cell lysate. Wild type HAP1 whole cell lysate.
General notes	<p>This product has switched from a hybridoma to recombinant production method on 02nd November 2020.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 7.3</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 0.05% BSA, 40% Glycerol (glycerin, glycerine), 59% PBS</p>
Purity	Protein A purified
Clonality	Monoclonal
Clone number	MT3.1
Isotype	IgG1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab39253 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★ (1)	Use at an assay dependent concentration. Predicted molecular weight: 22 kDa.
IHC-P	★★★★★ (1)	1/100. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Target

Function

Involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) in DNA. Repairs alkylated guanine in DNA by stoichiometrically transferring the alkyl group at the O-6 position to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.


Sequence similarities

Belongs to the MGMT family.

Cellular localization

Nucleus.

Images



All lanes : Anti-MGMT antibody [MT3.1] (ab39253) at 1/500 dilution

Lane 1 : Wild type HAP1 whole cell lysate


Lane 2 : MGMT knockout HAP1 whole cell lysate

Lane 3 : MCF7 (human breast adenocarcinoma epithelial cell)

Lane 4 : Jurkat (human T cell leukemia T lymphocyte)


Lysates/proteins at 20 µg per lane.

Predicted band size: 22 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGMT antibody [MT3.1] (ab39253)

Immunohistochemical analysis of paraffin-embedded human prostate carcinoma tissue labeling MGMT with ab39253 at 1/200 followed by a ready to use secondary. Positive staining on human prostate carcinoma. The section was incubated with ab39253 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use secondary. Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGMT antibody [MT3.1] (ab39253)

Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue labeling MGMT with ab39253 at 1/200 followed by a ready to use secondary. Positive staining on human lung carcinoma. The section was incubated with ab39253 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use secondary. Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.



All lanes : Anti-MGMT antibody [MT3.1] (ab39253) at 1/500 dilution

Lane 1 : MCF7 (human breast adenocarcinoma epithelial cell), whole cell lysate

Lane 2 : HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate

Lane 3 : Jurkat (human T cell leukemia T lymphocyte), whole cell lysate

Lane 4 : HCT116 (human colorectal carcinoma epithelial cell), whole cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Peroxidase-Conjugated Goat anti-Mouse IgG (H+L) at 1/10000 dilution

Predicted band size: 22 kDa

Observed band size: 22 kDa

Exposure time: 92 seconds




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGMT antibody [MT3.1] (ab39253)

This image is courtesy of an anonymous abreview.

ab39253 staining MGMT in human stomach tissue sections by Immunohistochemistry. Tissue was fixed with formaldehyde and a heat mediated antigen retrieval step was performed using 10mM citrate buffer pH6.0. Samples were then blocked with 5% serum for 1 hour at 23°C followed by incubation with the primary antibody, at a 1/100 dilution, for 1 hour at 23°C. An undiluted HRP-conjugated goat anti-mouse IgG polyclonal was used as secondary antibody.

This image was generated from the hybridoma version.



Western blot - Anti-MGMT antibody [MT3.1] (ab39253)

This image is courtesy of an anonymous Abreview

All lanes : Anti-MGMT antibody [MT3.1] (ab39253) at 1/500 dilution (for 16 hours at 4°C)

All lanes : Human OE33 cells - whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Rabbit Anti-Mouse IgG H&L (HRP) ([ab6728](#)) at 1/1000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 22 kDa

Observed band size: 21 kDa

Exposure time: 30 seconds

Blocking Step: 5% BSA for 1 hour at 23°C

This image was generated from the hybridoma version.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors