







Product datasheet

Anti-Myogenin antibody [F5D] - BSA and Azide free ab212667

1 Image

Overview

Product name	Anti-Myogenin antibody [F5D] - BSA and Azide free
Description	Mouse monoclonal [F5D] to Myogenin - BSA and Azide free
Host species	Mouse
Tested applications	Suitable for: IHC-P
Species reactivity	Reacts with: Human
Immunogen	<p>This product was produced with the following immunogens:</p> <p>Synthetic peptide corresponding to Rat Myogenin aa 50-150. Database link: P20428</p> <p>Recombinant fragment corresponding to Rat Myogenin aa 1-250. Database link: P20428</p> <div>  Run BLAST with  </div> <div>  Run BLAST with  </div> <div>  Run BLAST with  </div>
Positive control	Rh-30 cells; skeletal muscle or rhabdomyosarcoma tissues
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Constituent: 100% PBS
Carrier free	Yes

Purity	Protein G purified
Purification notes	Purified from bioreactor concentrate.
Clonality	Monoclonal
Clone number	F5D
Isotype	IgG1
Light chain type	kappa

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab212667 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use a concentration of 0.5 - 1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. for 30 minutes at room temperature.

Target

Function	Involved in muscle differentiation (myogenic factor). Induces fibroblasts to differentiate into myoblasts. Probable sequence specific DNA-binding protein.
Sequence similarities	Contains 1 basic helix-loop-helix (bHLH) domain.
Cellular localization	Nucleus.

Images



Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human rhabdomyosarcoma tissue section labeling Myogenin with ab212667 at 1 µg/mL.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Myogenin antibody [F5D]
- BSA and Azide free (ab212667)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors