


Product datasheet

Anti-PARK7/DJ1 antibody ab18257

KO VALIDATED

★★★★★ [11 Abreviews](#) [51 References](#) [5 Images](#)

Overview

Product name	Anti-PARK7/DJ1 antibody
Description	Rabbit polyclonal to PARK7/DJ1
Host species	Rabbit
Specificity	Replenishment batches of our polyclonal antibody, ab18257 are tested in WB. Previous batches were additionally validated in ICC/IF and IHC-P. These applications are still expected to work and are covered by our Abpromise guarantee. You may also be interested in our alternative recombinant antibody, ab76008 .
Tested applications	Suitable for: IHC-P, ICC/IF, WB
Species reactivity	Reacts with: Mouse, Rat, Human Predicted to work with: Rabbit, Cow, Dog 
Immunogen	Synthetic peptide corresponding to Human PARK7/DJ1 aa 150 to the C-terminus (C terminal) conjugated to keyhole limpet haemocyanin. (Peptide available as ab18659)
Positive control	WB: HeLa, Jurkat, PC12 and NIH 3T3 whole cell lysates, mouse brain, liver, heart, kidney, pancreas, testis, skeletal muscle, spinal cord and ovary tissue lysates and rat brain, liver, heart and kidney tissue lysates. IHC-P: Human parathyroid tissue. ICC/IF: HeLa cells.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

Storage buffer

pH: 7.40
Preservative: 0.02% Sodium azide
Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

Purity

Immunogen affinity purified

Clonality

Polyclonal

Isotype

IgG

Applications**The Abpromise guarantee**

Our **Abpromise guarantee** covers the use of ab18257 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration.
ICC/IF	★★★★★ (5)	Use a concentration of 1 µg/ml.
WB	★★★★★ (5)	Use a concentration of 1 µg/ml. Detects a band of approximately 24 kDa (predicted molecular weight: 20 kDa).

Target**Function**

Protects cells against oxidative stress and cell death. Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. May act as an atypical peroxiredoxin-like peroxidase that scavenges hydrogen peroxide. Following removal of a C-terminal peptide, displays protease activity and enhanced cytoprotective action against oxidative stress-induced apoptosis. Stabilizes NFE2L2 by preventing its association with KEAP1 and its subsequent ubiquitination. Binds to OTUD7B and inhibits its deubiquitinating activity. Enhances RELA nuclear translocation. Binds to a number of mRNAs containing multiple copies of GG or CC motifs and partially inhibits their translation but dissociates following oxidative stress. Required for correct mitochondrial morphology and function and for autophagy of dysfunctional mitochondria. Regulates astrocyte inflammatory responses. Acts as a positive regulator of androgen receptor-dependent transcription. Prevents aggregation of SNCA. Plays a role in fertilization. Has no proteolytic activity. Has cell-growth promoting activity and transforming activity. May function as a redox-sensitive chaperone.

Tissue specificity

Highly expressed in pancreas, kidney, skeletal muscle, liver, testis and heart. Detected at slightly lower levels in placenta and brain. Detected in astrocytes, Sertoli cells, spermatogonia, spermatids and spermatozoa.

Involvement in disease

Defects in PARK7 are the cause of Parkinson disease type 7 (PARK7) [MIM:606324]. A neurodegenerative disorder characterized by resting tremor, postural tremor, bradykinesia,

muscular rigidity, anxiety and psychotic episodes. PARK7 has onset before 40 years, slow progression and initial good response to levodopa. Some patients may show traits reminiscent of amyotrophic lateral sclerosis-parkinsonism/dementia complex (Guam disease).

Sequence similarities

Belongs to the peptidase C56 family.

Post-translational modifications

Sumoylated on Lys-130 by PIAS2 or PIAS4; which is enhanced after ultraviolet irradiation and essential for cell-growth promoting activity and transforming activity.


Cys-106 is easily oxidized to sulfinic acid.

Undergoes cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress.

Cellular localization

Cytoplasm. Nucleus. Mitochondrion. Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage. Detected in tau inclusions in brains from neurodegenerative disease patients.

Images



Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: PARK7/DJ1 knockout HAP1 cell lysate (20 µg)

Lane 3: HeLa cell lysate (20 µg)

Lane 4: Human brain tissue lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab18257 observed at 24 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab18257 was shown to specifically react with PARK/DJ1 in wild-type HAP1 cells. No band was observed when PARK/DJ1 knockout samples were used. Wild-type and PARK/DJ1 knockout samples were subjected to SDS-PAGE. ab18257 and [ab8245](#) (loading control to GAPDH) were diluted to 1 µg/mL and 1/10,000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10,000 dilution for 1 hr at room temperature before imaging.



All lanes : Anti-PARK7/DJ1 antibody (ab18257) at 1 µg/ml

Lane 1 : Brain (Mouse) Tissue Lysate

Lane 2 : Liver (Mouse) Tissue Lysate - normal tissue

Lane 3 : Heart (Mouse) Tissue Lysate

Lane 4 : Kidney (Mouse) Tissue Lysate

Lane 5 : Mouse pancreas tissue lysate - total protein ([ab29363](#))

Lane 6 : Testis (Mouse) Tissue Lysate - normal tissue

Lane 7 : Mouse skeletal muscle tissue lysate - total protein

([ab29711](#))

Lane 8 : Spinal Cord (Mouse) Tissue Lysate

Lane 9 : Ovary (Mouse) Tissue Lysate - normal tissue

Lane 10 : PC12 (Rat adrenal pheochromocytoma cell line) Whole Cell Lysate

Lane 11 : Brain (Rat) Tissue Lysate - normal tissue

Lane 12 : Liver (Rat) Tissue Lysate

Lane 13 : Heart (Rat) Tissue Lysate

Lane 14 : Kidney (Rat) Whole Cell Lysate - normal tissue

([ab29480](#))

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : IRDye 680 Conjugated Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 20 kDa

Observed band size: 24 kDa


Additional bands at: 15 kDa. We are unsure as to the identity of these extra bands.



Immunocytochemistry/ Immunofluorescence - Anti-PARK7/DJ1 antibody (ab18257)

ab18257 staining PARK7/DJ1 in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab18257 at 1µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Western blot - Anti-PARK7/DJ1 antibody (ab18257)

All lanes : Anti-PARK7/DJ1 antibody (ab18257) at 1 µg/ml

Lane 1 : Jurkat lysate

Lane 2 : HeLa lysate

Lane 3 : 3T3 lysate

Lane 4 : Jurkat lysate with Human PARK7/DJ1 peptide (**ab18659**) at 1 µg/ml

Lane 5 : HeLa lysate with Human PARK7/DJ1 peptide (**ab18659**) at 1 µg/ml

Lane 6 : 3T3 lysate with Human PARK7/DJ1 peptide (**ab18659**) at 1 µg/ml


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Alexa Fluor Goat polyclonal to Rabbit IgG (680) at 1/10000 dilution

Predicted band size: 20 kDa

Observed band size: 24 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors