


Product datasheet

PerCP/Cy5.5® Anti-CD4 antibody [GK1.5] ab210265

[1 Image](#)

Overview

Product name	PerCP/Cy5.5® Anti-CD4 antibody [GK1.5]
Description	PerCP/Cy5.5® Rat monoclonal [GK1.5] to CD4
Host species	Rat
Conjugation	PerCP/Cy5.5®. Ex: 482nm, Em: 690nm
Tested applications	Suitable for: Flow Cyt
Species reactivity	Reacts with: Mouse Predicted to work with: Syrian hamster 
Immunogen	Tissue, cells or virus corresponding to Mouse CD4. Mouse CTL clone V4.
Positive control	C57Bl/6 splenocytes.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Store In the Dark.
Storage buffer	<p>pH: 7.20</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: 0.12% Monobasic dihydrogen sodium phosphate, 0.1% Gelatin, 0.87% Sodium chloride</p>
Purity	Protein A purified
Purification notes	ab210265 was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.
Clonality	Monoclonal

Clone number	GK1.5
Isotype	IgG2b
Light chain type	kappa

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab210265 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use at an assay dependent concentration. Use 0.06 µg.

Target

Function	Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell activation. Induces the aggregation of lipid rafts.
Sequence similarities	Contains 3 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Post-translational modifications	Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.
Cellular localization	Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope polypeptide gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum.

Images



Flow cytometric analysis of C57Bl/6 splenocytes labeling CD4 with 0.06 µg ab210265 (solid line), compared with 0.06 µg Per-Cy5.5 Rat IgG2b isotype control (dashed line).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors