


Product datasheet

Anti-PIST antibody [EPR4079] ab109119

Recombinant RabMAb

[3 References](#) [9 Images](#)

Overview

Product name	Anti-PIST antibody [EPR4079]
Description	Rabbit monoclonal [EPR4079] to PIST
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, WB, IHC-P
Species reactivity	Reacts with: Human Predicted to work with: Mouse 
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: A375, 293T, and PC-3 cell lysates and human prostate and fetal kidney tissue lysates. IHC-P: Human stomach tissue. ICC/IF: HeLa cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	<p>pH: 7.20</p> <p>Preservative: 0.05% Sodium azide</p> <p>Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant</p>
Purity	Protein A purified

Clonality	Monoclonal
Clone number	EPR4079
Isotype	IgG

Applications


The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab109119 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/50 - 1/100.
WB		1/1000 - 1/10000. Predicted molecular weight: 51 kDa.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Target

Function	Plays a role in intracellular protein trafficking and degradation. May regulate CFTR chloride currents and acid-induced ACCN3 currents by modulating cell surface expression of both channels. May also regulate the intracellular trafficking of the ADR1B receptor. May play a role in autophagy. Overexpression results in CFTR intracellular retention and degradation in the lysosomes.
Tissue specificity	Ubiquitously expressed.
Involvement in disease	Note=A chromosomal aberration involving GOPC is found in a glioblastoma multiforme sample. An intra-chromosomal deletion del(6)(q21q21) is responsible for the formation of GOPC-ROS1 chimeric protein which has a constitutive receptor tyrosine kinase activity.
Sequence similarities	Contains 1 PDZ (DHR) domain.
Domain	The PDZ domain mediates interactions with FZD5, FZD8, ACCN3, GRID2, CLCN3 (By similarity). Mediates also interaction with CFTR and ADRB1. The coiled-coil region probably mediates association to membranes, targeting to the Golgi, and interactions with GOLGA3, and STX6. May also mediate interaction with RHOQ.
Cellular localization	Cytoplasm. Golgi apparatus membrane. Golgi apparatus > trans-Golgi network membrane. Cell junction > synapse. Cell junction > synapse > postsynaptic cell membrane > postsynaptic density. Cell projection > dendrite. Enriched in synaptosomal and postsynaptic densities (PSD) fractions (By similarity). Expressed in cell bodies and dendrites of Purkinje cells (By similarity). Localized at the trans-Golgi network (TGN) of spermatids and the medulla of round spermatides.

Images



All lanes : Anti-PIST antibody [EPR4079] (ab109119) at 1/1000 dilution

Lane 1 : A375 cell lysate

Lane 2 : 293T cell lysate


Lane 3 : Fetal kidney lysate

Lane 4 : PC-3 cell lysate


Lane 5 : Human prostate lysate

Lysates/proteins at 10 µg per lane.


Predicted band size: 51 kDa



ab109119 stained HeLa cells. The cells were 4% formaldehyde fixed for 10 minutes at room temperature and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1hour at room temperature to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab109119 at 1/100) overnight at +4°C. The secondary antibody (pseudo-colored green) was Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed (**ab150081**) used at a 1/1000 dilution for 1hour at room temperature. Alexa Fluor® 594 WGA was used to label plasma membranes (pseudo-colored red) at a 1/200 dilution for 1hour at room temperature. DAPI was used to stain the cell nuclei (pseudo-colored blue) at a concentration of 1.43µM for 1hour at room temperature.




Immunohistochemical staining of PIST expression in paraffin-embedded Human stomach tissue using ab109119 at 1/250 dilution. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PIST antibody [EPR4079] (ab109119)


ab109119 showing positive staining in Normal brain tissue.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PIST antibody [EPR4079] (ab109119)

ab109119 showing positive staining in Normal breast tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PIST antibody [EPR4079] (ab109119)

ab109119 showing positive staining in Normal kidney tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PIST antibody [EPR4079] (ab109119)

ab109119 showing positive staining in Normal liver tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PIST antibody [EPR4079] (ab109119)

ab109119 showing positive staining in Normal ovary tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-PIST antibody [EPR4079] (ab109119)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors