

Product datasheet

Anti-RKIP antibody [EPR2875Y] ab76582

Recombinant RabMAb

★★★★★ **5 Abreviews** **18 References** [10 Images](#)

Overview

Product name	Anti-RKIP antibody [EPR2875Y]
Description	Rabbit monoclonal [EPR2875Y] to RKIP
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), IHC-P, WB, ICC/IF Unsuitable for: IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide within Human RKIP aa 50-150. The exact sequence is proprietary.
Positive control	WB: HeLa, A549, HepG2, A431, mouse brain, mouse testis and rat brain cell lysates; IHC-P: human ovarian carcinoma and brain, mouse kidney, and rat testis tissues; ICC/IF: HepG2 cells; Flow Cyt (intra): HeLa cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR2875Y

Isotype

IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab76582 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/20.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .
WB	★★★★★ (3)	1/1000 - 1/5000. Detects a band of approximately 21 kDa (predicted molecular weight: 21 kDa).
ICC/IF	★★★★★ (2)	1/50 - 1/100.

Application notes

Is unsuitable for IP.

Target

Function

Binds ATP, opioids and phosphatidylethanolamine. Has lower affinity for phosphatidylinositol and phosphatidylcholine. Serine protease inhibitor which inhibits thrombin, neuropsin and chymotrypsin but not trypsin, tissue type plasminogen activator and elastase. HCNP may be involved in the function of the presynaptic cholinergic neurons of the central nervous system. HCNP increases the production of choline acetyltransferase but not acetylcholinesterase. Seems to be mediated by a specific receptor.


Sequence similarities

Belongs to the phosphatidylethanolamine-binding protein family.

Cellular localization

Cytoplasm.

Images



Western blot - Anti-RKIP antibody [EPR2875Y] (ab76582)

All lanes : Anti-RKIP antibody [EPR2875Y] (ab76582) at 1/1000 dilution (Purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : A549 (Human lung carcinoma epithelial cell) whole cell lysates

Lane 3 : Mouse brain lysates

Lane 4 : Mouse testis lysates

Lane 5 : Rat brain lysates


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution


Predicted band size: 21 kDa

Observed band size: 21 kDa




Immunocytochemistry/ Immunofluorescence - Anti-RKIP antibody [EPR2875Y] (ab76582)

Immunocytochemistry/ Immunofluorescence analysis of HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling RKIP with purified ab76582 at 1:50 dilution (2.6 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RKIP antibody
[EPR2875Y] (ab76582)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat testis tissue sections labeling RKIP with purified ab76582 at 1/500 dilution (0.26 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Flow Cytometry (Intracellular) - Anti-RKIP antibody
[EPR2875Y] (ab76582)

Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labeling RKIP (red) with purified ab76582 at a 1/20 dilution. Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. A goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal IgG (**ab172730**). Blue (unlabeled control) - Cells without incubation with the primary and secondary antibodies.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RKIP antibody
[EPR2875Y] (ab76582)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse kidney tissue sections labeling RKIP with purified ab76582 at 1/500 dilution (0.26 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RKIP antibody
[EPR2875Y] (ab76582)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human ovarian carcinoma tissue sections labeling RKIP with purified ab76582 at 1/500 dilution (0.26 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-RKIP antibody [EPR2875Y]
(ab76582)

All lanes : Anti-RKIP antibody [EPR2875Y] (ab76582) at 1/5000 dilution (unpurified)

Lane 1 : HeLa cell lysate

Lane 2 : A549 cell lysate

Lane 3 : HepG2 cell lysate

Lane 4 : A431 cell lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 21 kDa


Observed band size: 21 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-RKIP antibody
[EPR2875Y] (ab76582)

Staining of formalin-fixed, paraffin-embedded human brain tissue
using 1/250 ab76582 (unpurified).

Perform heat mediated antigen retrieval before commencing with
IHC staining protocol.



Western blot - Anti-RKIP antibody [EPR2875Y]
(ab76582)

Anti-RKIP antibody [EPR2875Y] (ab76582) at 1/5000 dilution
(unpurified) + Recombinant Human RKIP protein ([ab87892](#)) at 0.1
µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ([ab97080](#)) at
1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 21 kDa

Exposure time: 30 seconds

Why choose a recombinant antibody?



**Research with
confidence**
Consistent and
reproducible results



**Long-term and
scalable supply**
Recombinant
technology



**Success from the
first experiment**
Confirmed
specificity



**Ethical standards
compliant**
Animal-free
production

Anti-RKIP antibody [EPR2875Y] (ab76582)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors