abcam

Product datasheet

Anti-RRN3 (phospho S649) antibody ab138651

2 References 1 Image

Overview

Product name Anti-RRN3 (phospho S649) antibody

Description Rabbit polyclonal to RRN3 (phospho S649)

Host species Rabbit

Tested applications Suitable for: WB

Species reactivity Reacts with: Human

Immunogen Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

Positive control Extracts from starved Jurkat cells.

General notes

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at -20°C.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.88% Sodium chloride

PBS is without Mg²⁺ and Ca²⁺

Purity Immunogen affinity purified

Purification notes ab138651 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-

specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation.

Clonality Polyclonal

Isotype IgG

1

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab138651 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/1000. Predicted molecular weight: 74 kDa.

Target

Function Required for efficient transcription initiation by RNA polymerase I. Required for the formation of

the competent preinitiation complex (PIC). Dissociates from pol I as a consequence of transcription. In vitro, cannot activate transcription in a subsequent transcription reaction.

Sequence similarities Belongs to the RRN3 family.


Post-translational modifications

Cellular localization

Phosphorylation is required for participation in rDNA transcription.

Nucleus > nucleolus.

Images



Anti-RRN3 (phospho S649) antibody (ab138651) at 1/500 dilution + 24 hour starved Jurkat cell extract at 30 µg

Predicted band size: 74 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors