


Product datasheet

Anti-Scavenging Receptor SR-BI antibody [EP1556Y] ab52629

KO VALIDATED Recombinant RabMAB[®]

★★★★★ [4 Abreviews](#) [54 References](#) [8 Images](#)

Overview

Product name	Anti-Scavenging Receptor SR-BI antibody [EP1556Y]
Description	Rabbit monoclonal [EP1556Y] to Scavenging Receptor SR-BI
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P Unsuitable for: Flow Cyt or ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Human Predicted to work with: Sheep 
Immunogen	Synthetic peptide within Human Scavenging Receptor SR-BI aa 50-150 (N terminal). The exact sequence is proprietary. Database link: Q8WTV0
Positive control	WB: Mouse liver tissue lysate. IHC-P: Human liver tissue.
General notes	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production For more information see here . Our RabMAB [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAB[®] patents .

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

Purity	Protein A purified
Clonality	Monoclonal
Clone number	EP1556Y
Isotype	IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab52629 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★ (2)	1/1000 - 1/2000. Detects a band of approximately 80 kDa.
IHC-P		1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Application notes Is unsuitable for Flow Cyt or ICC/IF.

Target

Function Receptor for different ligands such as phospholipids, cholesterol ester, lipoproteins, phosphatidylserine and apoptotic cells. Probable receptor for HDL, located in particular region of the plasma membrane, called caveolae. Facilitates the flux of free and esterified cholesterol between the cell surface and extracellular donors and acceptors, such as HDL and to a lesser extent, apoB-containing lipoproteins and modified lipoproteins. Probably involved in the phagocytosis of apoptotic cells, via its phosphatidylserine binding activity. Receptor for hepatitis C virus glycoprotein E2. Binding between SCARB1 and E2 was found to be independent of the genotype of the viral isolate. Plays an important role in the uptake of HDL cholesteryl ester.


Tissue specificity Widely expressed.

Sequence similarities Belongs to the CD36 family.

Post-translational modifications N-glycosylated.

Cellular localization Cell membrane. Membrane > caveola. Predominantly localized to cholesterol and sphingomyelin-enriched domains within the plasma membrane, called caveolae.

Images



Western blot - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)

All lanes : Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : SCARB1 knockout HEK-293T cell lysate


Lane 3 : Human Liver cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Observed band size: 70,75 kDa

False colour image of Western blot: Anti-Scavenging Receptor SR-BI antibody [EP1556Y] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab52629 was shown to bind specifically to Scavenging Receptor SR-BI. A band was observed at 70/75 kDa in wild-type HEK-293T cell lysates with no signal observed at this size in SCARB1 knockout cell line [ab282646](#) (knockout cell lysate [ab283046](#)). To generate this image, wild-type and SCARB1 knockout HEK-293T cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Western blot - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: Scavenging Receptor SR-BI knockout HAP1 whole cell lysate (20 µg)

Lane 3: HepG2 whole cell lysate (20 µg)

Lane 4: Human liver whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab52629 observed at 80 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab52629 was shown to specifically react with Scavenging Receptor SR-BI in wild-type HAP1 cells as signal was lost in Scavenging Receptor SR-BI knockout cells. Wild-type and Scavenging Receptor SR-BI knockout samples were subjected to SDS-PAGE. Ab52629 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/2000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)

All lanes : Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629) at 1/1000 dilution (purified)

Lane 1 : Mouse liver lysate

Lane 2 : Rat liver lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Anti-rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Observed band size: 80 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



Western blot - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)

All lanes : Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629) at 1/1000 dilution (purified)

Lane 1 : Human fetal liver lysate

Lane 2 : HepG2 lysate

Lane 3 : PC-3 cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Anti-rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Observed band size: 80 kDa


Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



Immunohistochemical staining of paraffin embedded human liver with purified ab52629 at a working dilution of 1/500. The secondary antibody used is **ab97051**, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)




Western blot - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)

Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629) at 1/2000 dilution (unpurified) + Mouse liver lysate at 10 µg

Secondary

goat anti-rabbit HRP at 1/2000 dilution

Observed band size: 80 kDa



Western blot - Anti-Scavenging Receptor SR-BI antibody [EP1556Y] (ab52629)

THP1 cells were incubated at 37°C for 40h with vehicle control (0 µM) and different concentrations of sodium salicylate ([ab120746](#)). Decreased expression of scavenging receptor SR-BI in THP1 cells correlates with an increase in sodium salicylate concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 µg of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with unpurified ab52629 at 1/2000 dilution and [ab8227](#) at 1 µg/ml overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP ([ab97051](#)) at 1/10000 dilution and visualised using ECL development solution.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Scavenging Receptor SR-BI antibody
[EP1556Y] (ab52629)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors