

## Product datasheet

# Anti-SNX19 antibody ab121401

2 Images

### Overview

---

<b>Product name</b>	Anti-SNX19 antibody
<b>Description</b>	Rabbit polyclonal to SNX19
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> ICC/IF, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Human
<b>Immunogen</b>	antigen, corresponding to amino acids 625-750 of Human SNX19.
<b>Positive control</b>	Human kidney tissue.
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>Storage buffer</b>	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

### Applications

---

**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab121401 in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use a concentration of 0.25 - 2 µg/ml.
IHC-P		1/50 - 1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Perform Heat Induced Epitope Retrieval (HIER) at pH 6 using a pressure boiler as heat source.

## Target


<b>Function</b>	May be involved in several stages of intracellular trafficking.
<b>Sequence similarities</b>	Belongs to the sorting nexin family. Contains 1 PX (phox homology) domain. Contains 1 PXA domain.
<b>Domain</b>	The PX domain mediates specific binding to membranes enriched in phosphatidylinositol-3-phosphate (PtdIns(P3)).
<b>Cellular localization</b>	Cytoplasmic vesicle membrane.

## Images



Immunofluorescent staining of Human cell line U-2 OS shows positivity in mitochondria. Recommended concentration of ab121401 1-4 µg/ml. Cells treated with PFA/Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-SNX19 antibody (ab121401)



ab121401 at 1/75 dilution staining SNX19 in Paraffin-embedded Human Kidney tissue by Immunohistochemistry.

Note: strong cytoplasmic positivity in cells in tubules.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SNX19 antibody (ab121401)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors