

Product datasheet

Anti-TNF alpha antibody [52B83] ab1793

★★★★★ [18 Abreviews](#) [238 References](#) [4 Images](#)

Overview

Product name	Anti-TNF alpha antibody [52B83]
Description	Mouse monoclonal [52B83] to TNF alpha
Host species	Mouse
Specificity	This antibody detects both natural and recombinant TNFa. It does not cross-react with TNF beta or lymphotoxin. It reacts with free soluble (17 kDa) and membrane (26 kDa) human TNF-alpha. It does not react with receptor-bound TNF-alpha. Non-specific background staining is observed in connective tissues.
Tested applications	Suitable for: ELISA, IHC-P, ICC/IF, Flow Cyt (Intra), IHC-Fr, WB
Species reactivity	Reacts with: Mouse, Guinea pig, Human, Chimpanzee, Zebrafish, Cynomolgus monkey, Rhesus monkey, Apterionotus leptorhynchus
Immunogen	Full length native protein (purified) corresponding to Human TNF alpha.
Positive control	Flow Cyt (Intra): RAW 264.7 cells.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.02% Sodium azide Constituents: 0.1% BSA, PBS
Purity	Protein G purified
Clonality	Monoclonal
Clone number	52B83
Myeloma	unknown

Isotype	IgG1
Light chain type	unknown

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab1793 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ELISA		Use at an assay dependent concentration.
IHC-P	★★★★★ (5)	Use a concentration of 2.5 µg/ml.
ICC/IF	★★★★★ (2)	Use a concentration of 5 µg/ml.
Flow Cyt (Intra)		1/10. (methanol fixed cells) ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IHC-Fr	★★★★★ (3)	Use a concentration of 2 µg/ml. Fixed in acetone for 10 minutes
WB	★★★★★ (8)	Use a concentration of 2 µg/ml. Predicted molecular weight: 25 kDa. TNF-alpha is normally secreted as a homotrimer with a molecular mass of 52 kDa. Monomeric TNF-alpha is 17.4 kDa A reduced sample treatment and 15% SDS-Page was used.

Target


Function	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFB. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation.
Involvement in disease	Genetic variations in TNF are a cause of susceptibility psoriatic arthritis (PSORAS) [MIM:607507]. PSORAS is an inflammatory, seronegative arthritis associated with psoriasis. It is a heterogeneous disorder ranging from a mild, non-destructive disease to a severe, progressive, erosive arthropathy. Five types of psoriatic arthritis have been defined: asymmetrical oligoarthritis characterized by primary involvement of the small joints of the fingers or toes; asymmetrical arthritis which involves the joints of the extremities; symmetrical polyarthritis characterized by a rheumatoidlike pattern that can involve hands, wrists, ankles, and feet; arthritis mutilans, which is a rare but deforming and destructive condition; arthritis of the sacroiliac joints and spine (psoriatic spondylitis).
Sequence similarities	Belongs to the tumor necrosis factor family.
Post-translational modifications	The soluble form derives from the membrane form by proteolytic processing. The membrane form, but not the soluble form, is phosphorylated on serine residues.

Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1. O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid.

Cellular localization


Secreted and Cell membrane.

Images




ab1793 staining TNF α in RAW 264.7 cells treated with (R,S)-rolipram (**ab120029**), by ICC/IF. Decrease in TNF α expression correlates with increased concentration of (R,S)-rolipram, as described in literature.

The cells were incubated at 37°C for 24h in media containing different concentrations of **ab120029** ((R,S)-rolipram) in DMSO, fixed with 100% methanol for 5 minutes at -20°C and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab1793 (5 μ g/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-mouse polyclonal antibody (**ab96879**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.



Overlay histogram showing RAW 264.7 cells stained with ab1793 (red line). The cells were fixed with methanol (5 min) and incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab1793, 1/10 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (**ab91353**, 2 μ g/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a decreased signal in RAW 264.7 cells fixed with 4% paraformaldehyde (10 min) used under the same conditions.


Please note that Abcam do not have data for use of this antibody on non-fixed cells. We welcome any customer feedback.



Immunocytochemistry/ Immunofluorescence - Anti-TNF alpha antibody [52B83] (ab1793)

ab1793 staining TNF α in RAW 264.7 cells treated with (R)-(-)-rolipram (**ab120031**), by ICC/IF. Decrease in TNF α expression correlates with increased concentration of (R)-(-)-rolipram, as described in literature.

The cells were incubated at 37°C for 24h in media containing different concentrations of **ab120031** ((R)-(-)-rolipram) in DMSO, fixed with 100% methanol for 5 minutes at -20°C and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab1793 (5 μ g/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-mouse polyclonal antibody (**ab96879**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.



Immunocytochemistry/ Immunofluorescence - Anti-TNF alpha antibody [52B83] (ab1793)

ab1793 staining TNF α in RAW 264.7 cells treated with (S)-(+)-rolipram (**ab120030**), by ICC/IF. Decrease in TNF α expression correlates with increased concentration of (S)-(+)-rolipram, as described in literature.

The cells were incubated at 37°C for 24h in media containing different concentrations of **ab120030** ((S)-(+)-rolipram) in DMSO, fixed with 100% methanol for 5 minutes at -20°C and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab1793(5 μ g/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-mouse polyclonal antibody (**ab96879**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors