


Anti-TNFRSF19L antibody ab96220

[3 References](#) [2 Images](#)

Overview

Product name	Anti-TNFRSF19L antibody
Description	Rabbit polyclonal to TNFRSF19L
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat 
Immunogen	Recombinant fragment, corresponding to a region within amino acids 35-366 of Human TNFRSF19L (NP_116260)
Positive control	293T, A431, H1299, HepG2, MOLT4 and Raji cell lysate. HeLaS3 whole cell lysate. U87 xenograft tissue.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
Storage buffer	pH: 7.00 Preservative: 0.01% Thimerosal (merthiolate) Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab96220 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/3000. Predicted molecular weight: 46 kDa.
IHC-P		1/100 - 1/500.

Target

Function

Mediates activation of NF-kappa-B. May play a role in T-cell activation.

Tissue specificity

Highest levels are in spleen, lymph node, thymus, peripheral blood leukocytes, bone marrow and fetal liver. Very low levels in skeletal muscle, testis and colon. Not detected in brain, kidney and pancreas.

Sequence similarities

Belongs to the RELT family.
Contains 1 TNFR-Cys repeat.


Post-translational modifications

Phosphorylated in vitro by OXSR1.

Cellular localization

Cell membrane. Cytoplasm.


Images



Anti-TNFRSF19L antibody (ab96220) at 1/1000 dilution + HeLa S3 whole cell lysate at 30 µg

Predicted band size: 46 kDa

10% SDS PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TNFRSF19L antibody (ab96220)

ab96220 at 1/100 dilution staining TNFRSF19L in paraffin-embedded U87 xenograft sections by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors