


### Anti-VCIP 135 antibody ab97812

[2 Images](#)

#### Overview

<b>Product name</b>	Anti-VCIP 135 antibody
<b>Description</b>	Rabbit polyclonal to VCIP 135
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Mouse, Rat 
<b>Immunogen</b>	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 955 and 1217 of VCIP 135 (Q96JH7)
<b>Positive control</b>	H1299 whole cell lysate; 293T, A431, HeLaS3, HepG2, MOLT4 or Raji lysates. SW480 xenograft (IHC-P)
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.00 Preservative: 0.01% Thimerosal (merthiolate) Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)
<b>Purity</b>	Immunogen affinity purified
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

#### Applications

## The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab97812 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/5000. Predicted molecular weight: 134 kDa.
IHC-P		Use at an assay dependent concentration.

## Target

### Function

Acts as a deubiquitinating enzyme. Necessary for VCP-mediated reassembly of Golgi stacks after mitosis. May play a role in VCP-mediated formation of transitional endoplasmic reticulum (tER). Mediates dissociation of the ternary complex containing STX5A, NSFL1C and VCP (By similarity). Hydrolyzes 'Lys-11'- and 'Lys-48'-linked polyubiquitin chains.


### Sequence similarities

Contains 1 OTU domain.

### Cellular localization

Endoplasmic reticulum. Golgi apparatus, Golgi stack. Associated with Golgi stacks and endoplasmic reticulum.


## Images



Anti-VCIP 135 antibody (ab97812) at 1/5000 dilution + H1299 whole cells lysate at 30 µg

**Predicted band size:** 134 kDa

7.5% SDS Page



Immunohistochemical analysis of paraffin-embedded SW480 xenograft, using ab97812 at 1/100 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-VCIP 135 antibody (ab97812)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors