


## Product datasheet

# Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] ab14734

KO VALIDATED

★★★★★ [52 Abreviews](#) [512 References](#) [12 Images](#)

### Overview

<b>Product name</b>	Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2]
<b>Description</b>	Mouse monoclonal [20B12AF2] to VDAC1/Porin + VDAC3
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> WB, ICC/IF, Flow Cyt
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Cow, Human <b>Predicted to work with:</b> Sheep, Goat, Cat, Dog, Pig, Drosophila melanogaster, Fish, Quail, Common marmoset, Dogfish, Catshark 
<b>Immunogen</b>	Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.
<b>Positive control</b>	WB: Isolated mitochondria from human, cow, rat and mouse heart. HepG2 cell lysate. ICC/IF: HeLa cells. Human fibroblasts. Flow Cyt: HepG2 cells.
<b>General notes</b>	<p>This antibody clone [20B12AF2] is manufactured by Abcam.</p> <p>If you require this antibody in a different buffer formulation or a different conjugate for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a> or you can find further information <a href="#">here</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C. Do Not Freeze.
<b>Storage buffer</b>	<p>pH: 7.50</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 0.36% HEPES, 0.88% Sodium chloride</p>

<b>Purity</b>	IgG fraction
<b>Purification notes</b>	Near homogeneity as judged by SDS-PAGE. The antibody was produced in vitro using hybridomas grown in serum-free medium, and then purified by biochemical fractionation.
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	20B12AF2
<b>Isotype</b>	IgG2b
<b>Light chain type</b>	kappa

## Applications


**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab14734 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
<b>WB</b>	★★★★★ (18)	Use a concentration of 1 µg/ml. Detects a band of approximately 39 kDa.
<b>ICC/IF</b>	★★★★★ (4)	Use at an assay dependent concentration.
<b>Flow Cyt</b>		Use 1µg for 10 <sup>6</sup> cells. <b>ab170192</b> - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.

## Target

**Cellular localization** VDAC1/Porin: Mitochondrion outer membrane. Cell membrane. VDAC3: Mitochondrial outer membrane.

## Images



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

**All lanes :** Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734) at 1/1000 dilution

**Lane 1 :** Wild-type Hap1 cell lysate

**Lane 2 :** VDAC1 knockout Hap1 cell lysate


**Lane 3 :** Wild-type HEK-293T cell lysate

**Lane 4 :** VDAC1 knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

**Lanes 1 -4:** Merged signal (red and green). Green - ab14734 observed at 31 kDa. Red - loading control, **ab181602** observed at 37 kDa. The lower band very close to VDAC1 is likely to be VDAC3.

ab14734 was shown to react with VDAC1/Porin + VDAC3 in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line **ab255444** (knockout cell lysate **ab263839**) was used. Wild-type and VDAC1 / Porin knockout samples were subjected to SDS-PAGE. ab14734 and Anti-GAPDH antibody [EPR16891] - Loading Control (**ab181602**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

This image is courtesy of an Abreview by Michiel Kroels.

ab14734 staining VDAC1/Porin + VDAC3 in human HeLa (Human epithelial cell line from cervix adenocarcinoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence).

Cells were fixed with paraformaldehyde, permeabilized with 0.1% Triton X-100 for 3 minutes and blocked with 0.2% serum for 60 minutes at 22°C. Samples were incubated with primary antibody (1/200 in 0.5% BSA and 0.02% Triton X100 in PBS) for 16 hours at 4°C. An FITC-conjugated goat anti-mouse IgG polyclonal was used as the secondary antibody at a dilution of 1/200.



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

**All lanes :** Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734) at 1/1000 dilution

**Lane 1 :** N-GST tagged Recombinant Human VDAC1 (aa1 to 283) protein with NFDM/TBST

**Lane 2 :** N-GST tagged Recombinant Human VDAC2 (aa 1 to 294) protein with NFDM/TBST

**Lane 3 :** N-GST tagged Recombinant Human VDAC3 (aa 1 to 283) protein with NFDM/TBST

Lysates/proteins at 0.01 µg per lane.

Blocking peptides at 5 % per lane.


**Observed band size:** 33 kDa

**Exposure time:** 10 seconds

Diluting buffer: 5% NFDM /TBST

N-GST tagged Recombinant Human VDAC1 protein is available as **ab132481**

N-GST tagged Recombinant Human VDAC2 protein is available as **ab152793**



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

**All lanes :** Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734) at 1/1000 dilution

**Lane 1 :** Wild type HAP1 whole cell lysate


**Lane 2 :** VDAC1 knockout HAP1 whole cell lysate

Lysates/proteins at 20 µg per lane.

**Observed band size:** 33 kDa

**Exposure time:** 5 seconds

Blocking and dilution buffer: 5% NFDM /TBST.



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

**All lanes :** Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734) at 1/1000 dilution

**Lane 1 :** HEK-293 (Human embryonic kidney epithelial cell) whole cell lysate with NFDM/TBST

**Lane 2 :** HAP1 (Wildtype control Human chronic myelogenous leukemia near-haploid cell line) whole cell lysate with NFDM/TBST

**Lane 3 :** MEF (Mus musculus Embryo Fibroblast) whole cell lysate with NFDM/TBST

**Lane 4 :** Rat brain tissue lysate with NFDM/TBST


Lysates/proteins at 20 µg per lane.

Blocking peptides at 5 % per lane.

**Observed band size:** 33 kDa

**Exposure time:** 5 seconds

Diluting buffer: 5% NFDM /TBST



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

Sun Y et al Voltage-dependent anion channels (VDACs) recruit Parkin to defective mitochondria to promote mitochondrial autophagy. J Biol Chem. 2012 Nov 23;287(48):40652-60. doi: 10.1074/jbc.M112.419721.


**(A) The Anti-VDAC1/Porin + VDAC3 antibody recognizes VDAC1 and VDAC3 but not VDAC2.**

Lysates from wild-type MEFs and VDAC1/3-/- MEFs were analyzed by western blotting with Anti-VDAC1/Porin + VDAC3 (top panel) or anti Tom20 antibodies (bottom panel).

**(B) The Anti-VDAC1/Porin + VDAC3 antibody does not recognize VDAC2 in VDAC1/3-/- MEFs.**

HA-tagged VDAC1, 2 or 3 were expressed in VDAC1/3-/- MEFs, immunoprecipitated with anti-HA antibodies and analyzed by western blotting with anti-HA antibodies (top panel) or Anti-VDAC1/Porin + VDAC3 antibodies (bottom panel). The Anti-VDAC1/Porin + VDAC3 antibodies recognize HA-VDAC1 and HA-VDAC3, but not HA-VDAC2 . Note that HA-VDAC3 migrates at a lower molecular weight than HA-VDAC1 and HA-VDAC2 .

Molecular weights are indicated in kDa.



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)


Li et al PLoS One. 2016 Feb 22;11(2):e0149728. doi: 10.1371/journal.pone.0149728. eCollection 2016. Fig 5.

Immunoblot analysis of endogenous VDAC1/Porin and VDAC3 in subcellular fractions of MA-10 cell lysates. VDAC1/Porin and VDAC3 are used as a mitochondrial marker proteins.

**Lane 1:** Whole cell lysate

**Lane 2:** Cytosolic fraction

**Lane 3:** Mitochondrial-enriched fraction



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

**All lanes :** Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

**Lane 1 :** Isolated mitochondria from human heart at 15 µg

**Lane 2 :** Isolated mitochondria from bovine heart at 6 µg


**Lane 3 :** Isolated mitochondria from rat heart at 30 µg

**Lane 4 :** Isolated mitochondria from mouse heart at 30 µg

**Lane 5 :** HepG2 (Human liver hepatocellular carcinoma cell line) cell lysate at 30 µg

**Observed band size:** 37 kDa

Extra bands in the mouse sample (lane 4) are due to the reaction of the secondary antibody with residual mouse blood in the heart tissue, as it is very difficult to entirely remove the blood from these small organs.




Flow Cytometry - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

Overlay histogram showing HepG2 (Human liver hepatocellular carcinoma cell line) cells stained with ab14734 (red line).

The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab14734, 1  $\mu$ g/ $1 \times 10^6$  cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2b [PLPV219] ([ab91366](#), 2  $\mu$ g/ $1 \times 10^6$  cells) used under the same conditions.

Acquisition of >5,000 events was performed.


This antibody gave a positive signal in HepG2 cells fixed with 80% methanol (5 min)/permeabilized in 0.1% PBS-Tween used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

Immunofluorescence using ab14734 at 0.2  $\mu$ g/ml on human fibroblasts (red).

Nuclei were labeled with DAPI (blue).



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

Image courtesy of an anonymous abreview.

Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734) at 1/500 dilution + MCF7 (Human breast adenocarcinoma cell line) cell lysate at 10 µg

#### Secondary


HRP-conjugated goat anti-mouse polyclonal at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Exposure time:** 3 minutes

0.5% TBS-tween + Lait 5% NaN<sub>3</sub> for 16 hours at 4°C.



Western blot - Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734)

This image is courtesy of an anonymous Abreview

**All lanes :** Anti-VDAC1/Porin + VDAC3 antibody [20B12AF2] (ab14734) at 1/5000 dilution

**Lanes 1-2 :** Rat brain cell lysate (homogenate)

**Lanes 3-4 :** Rat brain cell lysate (mitochondrial)

Lysates/proteins at 30 µg per lane.

#### Secondary

**All lanes :** HRP conjugated sheep anti-mouse IgG

Developed using the ECL technique.

Performed under reducing conditions.

**Observed band size:** 39 kDa

**Exposure time:** 5 minutes

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours



- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors