

Anti-ZMIZ2 antibody ab113277

[1 Image](#)

Overview

Product name	Anti-ZMIZ2 antibody
Description	Rabbit polyclonal to ZMIZ2
Host species	Rabbit
Specificity	ab113277 is predicted to not cross-react with other ZMIZ protein family members.
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Mouse
Immunogen	A 17 amino acid synthetic peptide corresponding to internal sequence amino acids of Human ZMIZ2.
Positive control	A20 cell lysate.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at 4°C (stable for up to 12 months).
Storage buffer	<p>pH: 7.2</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: 99% PBS</p>
Purity	Immunogen affinity purified
Purification notes	ab113277 is purified by affinity chromatography via a peptide column.
Clonality	Polyclonal
Isotype	IgG

Applications

Applications

The Abpromise guarantee

Our **Abpromise guarantee** covers the use of ab113277 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use a concentration of 0.25 - 0.5 µg/ml. Predicted molecular weight: 97 kDa.

Target

Function

Increases ligand-dependent transcriptional activity of AR and other nuclear hormone receptors.

Tissue specificity

Expressed most abundantly in testis with lower levels in heart, brain, pancreas, prostate and ovary.

Sequence similarities

Contains 1 SP-RING-type zinc finger.


Domain

The C-terminal proline-rich domain possesses a significant intrinsic transcriptional activity. This activity is inhibited by the N-terminus in the full-length protein.

Cellular localization

Nucleus. Detected at replication foci throughout S phase.

Images



Lane 1 : Anti-ZMIZ2 antibody (ab113277) at 0.25 µg/ml

Lane 2 : Anti-ZMIZ2 antibody (ab113277) at 0.5 µg/ml

All lanes : A20 cell lysate

Lysates/proteins at 15 µg per lane.

Predicted band size: 97 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors