

Recombinant human ALDH3A1 protein ab93463

1 Image

Description

Product name	Recombinant human ALDH3A1 protein
Biological activity	Specific activity is >5 units/ml and was obtained by measuring the increase of NADP in absorbance at 340 nm resulting from the reduction of NAD. One unit will oxidize 1.0 umole of acetaldehyde to acetic acid per minute at pH 8.0 at 25C in the presence of beta-NAD, potassium and thiols. Specific activity is >5 units/ml and was obtained by measuring the increase of NADP in absorbance at 340 nm resulting from the reduction of NAD. One unit will oxidize 1.0 umole of acetaldehyde to acetic acid per minute at pH 8.0 at 25C in the presence of beta-NAD, potassium and thiols.
Purity	> 95 % SDS-PAGE. ab93463 is purified using conventional chromatography techniques.
Expression system	Escherichia coli
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Human
Sequence	

MGSSHHHHHH SSGLVPRGSH MSKISEAVKR
ARAAFSSGRT RPLQFRIQQL EALQRLIQEQ
EQELVGALAA DLHKNEWNAY YEEVVVLEE
IEYMIQKLPE WAADEPVEKT PQTQQDELYI HSEPLGVVLV
IGTWNYPFNL TIQPMVGAIA AGNAVVLKPS ELSENMASLL
ATIIPQYLDK DLYPVINGGV PETTELLKER FDHILYTGST
GVGKIIMTAA AKHLTPVTLE LGGKSPCYVD
KNCDLDVACR RIAWGKFMNS GQTCVAPDYI
LCDPSIQNQI VEKLKSLKE FYGEDAKKSR DYGRISARH
FQRVMGLIEG QKVAYGGTGD AATRYIPTI LTDVDPQSPV
MQEEIFGPVL PIVCVRSLEE AIQFINQREK PLALYMFSSN
DKVIKKMAE TSSGGVAAND VMHITLHSL PFGGVGNSGM
GSYHGKKSFE TFSHRRSCLV RPLMNDEGLK
VRYPPSPAKM TQH

Specifications

Our **Abpromise guarantee** covers the use of **ab93463** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE
	Functional Studies
Form	Liquid


Preparation and Storage

Stability and Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
	pH: 8.00
	Constituents: 0.316% Tris HCl, 10% Glycerol (glycerin, glycerine), 0.58% Sodium chloride
	This product is an active protein and may elicit a biological response in vivo, handle with caution.

General Info

Function	ALDHs play a major role in the detoxification of alcohol-derived acetaldehyde. They are involved in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation. This protein preferentially oxidizes aromatic aldehyde substrates. It may play a role in the oxidation of toxic aldehydes.
Tissue specificity	High levels in stomach, esophagus and lung; low level in the liver and kidney.
Sequence similarities	Belongs to the aldehyde dehydrogenase family.
Cellular localization	Cytoplasm.

Images



15% SDS-PAGE analysis of ab93463 (3µg).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors