abcam

Product datasheet

Recombinant human CDK4 + CCND3 protein ab85646

1 References 5 Images

Description

Product name Recombinant human CDK4 + CCND3 protein

Biological activity The Specific activity of ab85646 was determined to be 16 nmol/min/mg.

Purity > 80 % Densitometry.

Affinity purified.

Expression system Baculovirus infected Sf9 cells

Protein length Full length protein

Animal free No

Nature Recombinant

Species Human

Specifications

Our Abpromise guarantee covers the use of ab85646 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications Western blot

Functional Studies

SDS-PAGE

Form Liquid

Additional notes ab56270 (Human Rb protein fragment) can be utilized as a substrate for assessing Kinase

activity

Preparation and Storage

Stability and Storage Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.

pH: 7.50

Constituents: 0.0038% EGTA, 0.00174% PMSF, 0.00385% DTT, 0.79% Tris HCl, 0.00292%

EDTA, 25% Glycerol (glycerin, glycerine), 0.87% Sodium chloride

This product is an active protein and may elicit a biological response in vivo, handle with caution.

. . .

1


Relevance

CDK4 is a member of the Ser/Thr protein kinase family. It is highly similar to the gene products of S. cerevisiae cdc28, and S. pombe cdc2. CDK4 is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is restricted to the G1/S phase, which is controlled by the regulatory subunits D type cyclins and CDK inhibitor p16(INK4a). CDK4 was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as its related proteins including D type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. CCND3 (Cyclin D3) belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. CCND3 forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. It has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb.


Cellular localization

CCND3: Cytoplasm. Nucleus.

Images




The specific activity of CDK4 + CCND3 (ab85646) was determined to be 13 nmol/min/mg as per activity assay protocol




SDS PAGE analysis of ab85646

SDS-PAGE - Recombinant human CDK4 + CCND3 protein (ab85646)




SDS PAGE analysis of ab85646





The Specific activity of ab85646 was determined to be 16 nmol/min/mg.





protein (ab85646)

SDS-PAGE showing ab85646 at approximately 58kDa (both proteins). $\label{eq:sdecomposition}$

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors