

Product datasheet

Recombinant Human Cytokeratin 8 protein
(denatured) ab156729

1 Image

Description	
Product name	Recombinant Human Cytokeratin 8 protein (denatured)
Purity	> 90 % SDS-PAGE.
Expression system	Escherichia coli
Accession	<u>P05787</u>
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Human
Sequence	MGSSHHHHHH SSGLVPRGSH MGSMsIRVTQ KSYKVSTSGP RAFSSRSYTS GPGSRISsss FSRVGSSNFR GGLGGGYGGA SGMGGITAVT VNQSLLSPLV LEVDPNIQAV RTQEKEQIKT LNNKFASFID KVRFLEQQNK MLETKWSLLQ QQKTARsNMD NMFESYINNl RRQLETlGQE KlKLEAELGN MQGLVEDFKN KYEDEINKRT EMENEfVLIK KDVDEAYMNK VELESrLEGL TDEINflRQL YEEEIRELQS QISDTsVVLs MDNSRSLDMD SIAEVKAQY EDIANRSRAE AESMYQIKYE ELQSLAGKHG DDLRRtKTEI SEMNrnISRL QAEIEGLKGQ RASLEAAIAD AEQRGELAiK DANAKLSELE AALQRAKQDM ARQLREYQEL MNVKLALDIE IATYRKlLEG EESRLESGMQ NMSIHTKtTS GYAGGLSSAY GGLTSPGLSY SLGSSFGSGA GSSsFSRTSS SRAVVVKKIE TRDGKLvSES SDVLpK
Predicted molecular weight	56 kDa including tags
Amino acids	1 to 483
Tags	His tag N-Terminus

Specifications

Our **Abpromise guarantee** covers the use of **ab156729** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications SDS-PAGE

Form Liquid

Preparation and Storage

Stability and Storage Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 2.4% Urea, 0.32% Tris HCl, 10% Glycerol (glycerin, glycerine)

General Info

Function Together with KRT19, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.

Tissue specificity Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma membrane in structures that contain dystrophin and spectrin. Expressed in gingival mucosa and hard palate of the oral cavity.


Involvement in disease Cirrhosis

Sequence similarities Belongs to the intermediate filament family.

Post-translational modifications Phosphorylation on serine residues is enhanced during EGF stimulation and mitosis. Ser-74 phosphorylation plays an important role in keratin filament reorganization. O-glycosylated. O-GlcNAcylation at multiple sites increases solubility, and decreases stability by inducing proteasomal degradation. O-glycosylated (O-GlcNAcylation), in a cell cycle-dependent manner.

Cellular localization Cytoplasm. Nucleus, nucleoplasm. Nucleus matrix.

Images



15% SDS-PAGE analysis of ab156729 (3µg).

SDS-PAGE - Recombinant Human Cytokeratin 8
protein (denatured) (ab156729)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors