

Recombinant Human UFL1 protein ab161615

1 Image

Description	
Product name	Recombinant Human UFL1 protein
Expression system	Wheat germ
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Human
Sequence	MADAWEEIRRLAADFQRAQFAEATQRLSERNCIEV NKLIA QKQLEV VHT LDGKEYTPAQISKEMRDELHVRGGRVNIVDLQQVINVDLIHI ENRIGDI IKSEKHVQLVLGQLIDENYLDRLAEEVNDKLQESGQVTISE LCKTYDLP G NFLTQALTQRLGRIISGHIDLDNRGVIFTEAFVARHKARIRGL FSAITRP TAVNSLISKYGFQEQLLYSVLEELVNSGRLRGTVVGGRQD KAVFVPDIYS RTQSTWVDSFFRQNGYLEFDALSRLGIPDAVSYKKRYKTT QLLFLKAAC VGQGLVDQVEASVEEAISSGTWVDIAPLLPTSLSVEDAAIL LQQVMRAFS KQASTVVFSDTV VVSEKFINDCTELFREL MHQKA EKEMK NNPVHLITEED LKQISTLESVSTSKDKKDERRRKATEGSGSMRGGGGGN AREYKIKK VKK KGRKDDSDDESQSSHTGKKKPEISFMFQDEIEDFLRKHI QDAPEEFISE LAEYLIKQV
Amino acids	1 to 509
Tags	GST tag N-Terminus

Specifications

Our **Abpromise guarantee** covers the use of **ab161615** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	Western blot ELISA
Form	Liquid
Additional notes	This product was previously labelled as K0776.


Preparation and Storage

Stability and Storage	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 8.00 Constituents: 0.31% Glutathione, 0.79% Tris HCl
------------------------------	--

General Info

Function	E3 UFM1-protein ligase that mediates ufmylation of target proteins such as DDRGK1/C20orf116. The function of ufmylation is unknown. May act as a tumor suppressor by inhibiting cell invasion, blocking NF-kappa-B signaling and increasing stability of CDK5RAP3.
Tissue specificity	Ubiquitously expressed, with a high expression in liver (at protein level). Low expression in several invasive hepatocellular carcinomas, such HepG2, Hep3B, HLE and PLC.
Sequence similarities	Belongs to the UFL1 family.
Post-translational modifications	Ubiquitinated, leading to its degradation by the proteasome. Interaction with CDK5RAP3 protects each other against ubiquitination and degradation via the proteasome.
Cellular localization	Endoplasmic reticulum. Cytoplasm. In cytoplasm colocalizes with CDK5RAP3.

Images



ab161615 on a 12.5% SDS-PAGE stained with Coomassie Blue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors