

Product datasheet

Recombinant mouse Macrophage Inflammatory Protein 3 alpha ab72992

[1 Image](#)

Description

Product name	Recombinant mouse Macrophage Inflammatory Protein 3 alpha
Biological activity	Biological activity is determined by its ability to chemoattract murine CCR6 transfected HEK/293 cells using a concentration range of 0.1-10.0 ng/ml.
Purity	> 95 % SDS-PAGE. Purity is greater than 95% as determined by reducing and non-reducing SDS-PAGE.
Expression system	Escherichia coli
Protein length	Full length protein
Animal free	No
Nature	Recombinant
Species	Mouse
Sequence	ASNYDCCLSY IQTPLPSRAI VGFTQMADE ACDINAIIFH TKKRKSVCAD PKQNWVKRAV NLLSLRVKKM

Specifications

Our **Abpromise guarantee** covers the use of **ab72992** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE Functional Studies
Form	Lyophilized

Preparation and Storage

Stability and Storage	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C. This product is an active protein and may elicit a biological response in vivo, handle with caution.
Reconstitution	Reconstitute with sterile water at a concentration of 0.1 - 0.5 mg/ml, which can be further diluted into other aqueous solutions. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

General Info

Function

Chemotactic factor that attracts lymphocytes and, slightly, neutrophils, but not monocytes. Inhibits proliferation of myeloid progenitors in colony formation assays. May be involved in formation and function of the mucosal lymphoid tissues by attracting lymphocytes and dendritic cells towards epithelial cells. C-terminal processed forms have been shown to be equally chemotactically active for leukocytes. Possesses antibacterial activity E.coli ATCC 25922 and S.aureus ATCC 29213.

Tissue specificity

Expressed predominantly in the liver, lymph nodes, appendix, peripheral blood lymphocytes, and fetal lung. Low levels seen in thymus, prostate, testis, small intestine and colon.

Sequence similarities

Belongs to the intercrine beta (chemokine CC) family.


Post-translational modifications

C-terminal processed forms which lack 1, 3 or 6 amino acids are produced by proteolytic cleavage after secretion from peripheral blood monocytes.

Cellular localization

Secreted.

Images



ab72992 used in Functional Studies.

Functional Studies - Recombinant mouse

Macrophage Inflammatory Protein 3 alpha (ab72992)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors