abcam

Product datasheet

Annexin V/ANXA5-Cy5 Apoptosis Staining / Detection Reagent ab14147

2 References

Overview

Product name Annexin V/ANXA5-Cy5 Apoptosis Staining / Detection Reagent

Tested applications Suitable for: FM, Flow Cyt

General notes This product is manufactured by BioVision, an Abcam company and was previously called 1013

Annexin V-Cy5 Reagent.

Annexin V/ANXA5-Cy5 Apoptosis Detection Reagent ab14147 can be used in a 10 min, onestep staining procedure to detect apoptosis by staining phosphatidylserine molecules which have translocated to the outside of the cell membrane. Analysis is by flow cytometry or fluorescence

microscopy.

This Annexin V/ANXA5-Cy5 reagent can be used with Annexin V/ANXA5 binding buffer <u>ab14084</u> (see protocol section for instructions). The Annexin V/ANXA5-Cy5 reagent is also available as

part of Annexin V/ANXA5-Cy5 apoptosis kit ab14150.

For more apoptosis assays, review the full set of **Annexin V/ANXA5 assays**, or the **apoptosis**

assay and apoptosis marker guide.

Properties

Form Liquid

Storage instructions Store at +4°C.

Storage buffer pH: 7.4

Preservative: 0.01% Thimerosal (merthiolate)

Constituents: PBS, 1% BSA

Function This protein is an anticoagulant protein that acts as an indirect inhibitor of the thromboplastin-

specific complex, which is involved in the blood coagulation cascade.

Involvement in disease Pregnancy loss, recurrent, 3

Sequence similarities Belongs to the annexin family.

Contains 4 annexin repeats.

DomainThe [IL]-x-C-x-x-[DE] motif is a proposed target motif for cysteine S-nitrosylation mediated by the

iNOS-S100A8/A9 transnitrosylase complex.

A pair of annexin repeats may form one binding site for calcium and phospholipid.

Post-translational S-nitrosylation is induced by interferon-gamma and oxidatively-modified low-densitity lipoprotein

1

Applications

The Abpromise guarantee Our Abpromise guarantee covers the use of ab14147 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
FM		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors