abcam

Product datasheet

Mouse brain tissue lysate - total protein (14 days) ab7189

| Product nameMouse brain tissue lysate - total protein (14 days)General notesSource: Outbred NIH-Swiss Albino mice.The mouse brain tissue was collected from fourteen (14) day-old mice. The lysate was prepa by homogenization in modified RIPA buffer (50 mM Tris -HCl, pH 7.4, 1% Triton X-100, 0.2% sodium deoxycholate, 0.2% sodium dodecylsulfate (SDS), 1 mM sodium ethylenediaminetetraacetate, 1 mM phenylmethylsulfonyl flouride, 5 µg/ml of aprotinin, 5 µg/m leupeptin). Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The tissue lysate was boiled for 5 min in 1 x SDS sa buffer (0.045 M Tris-HCl pH 6.8, 10% glycerol, 1% sodium dodecylsulfate, 0.01% bromopher blue), containing 0.05 M DTT. |
|---|
| The mouse brain tissue was collected from fourteen (14) day-old mice. The lysate was prepa by homogenization in modified RIPA buffer (50 mM Tris -HCI, pH 7.4, 1% Triton X-100, 0.2% sodium deoxycholate, 0.2% sodium dodecylsulfate (SDS), 1 mM sodium ethylenediaminetetraacetate, 1 mM phenylmethylsulfonyl flouride, 5 µg/ml of aprotinin, 5 µg/m leupeptin). Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The tissue lysate was boiled for 5 min in 1 x SDS sa buffer (0.045 M Tris-HCI pH 6.8, 10% glycerol, 1% sodium dodecylsulfate, 0.01% bromopher blue), containing 0.05 M DTT. |
| by homogenization in modified RIPA buffer (50 mM Tris -HCI, pH 7.4, 1% Triton X-100, 0.2% sodium deoxycholate, 0.2% sodium dodecylsulfate (SDS), 1 mM sodium ethylenediaminetetraacetate, 1 mM phenylmethylsulfonyl flouride, 5 µg/ml of aprotinin, 5 µg/m leupeptin). Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The tissue lysate was boiled for 5 min in 1 x SDS sa buffer (0.045 M Tris-HCI pH 6.8, 10% glycerol, 1% sodium dodecylsulfate, 0.01% bromopher blue), containing 0.05 M DTT. |
| |
| Tested applications Suitable for: WB |
| Properties |
| Mycoplasma free Yes |
| Form Liquid |
| Storage instructions Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. |
| Storage buffer pH: 7.20 Constituent: 100% SDS Sample Buffer |
| |

Applications

The Abpromise guarantee Our <u>Abpromise guarantee</u> covers the use of ab7189 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| WB | | Use at an assay dependent dilution. 10 μ g to 20 μ g per lane is recommended for mini gel. |

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.com/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors