

Product datasheet

Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] ab179463

Recombinant RabMAb

[275 References](#) [12 Images](#)

Overview

Product name	Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798]
Description	Rabbit monoclonal [EPR16798] to AKT1 + AKT2 + AKT3
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P, ICC/IF, IP, Flow Cyt (Intra)
Species reactivity	Reacts with: Mouse, Rat, Human, Xenopus laevis, Xenopus tropicalis
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: MCF7, HeLa, Hep G2 and A549 whole cell lysates; Human fetal brain, heart and kidney lysates; Mouse and Rat brain, heart, kidney and spleen lysates; Xenopus muscle lysate; AKT2 and AKT3 recombinant proteins. IHC-P: Human kidney, Mouse and Rat cerebral cortex. ICC/IF: K562 cells. Flow: A549 cells. IP: MCF7 whole cell lysate
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal

Clone number EPR16798
 Isotype IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab179463 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/10000. Detects a band of approximately 56 kDa (predicted molecular weight: 56 kDa).
IHC-P		1/250. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/100.
IP		1/100.
Flow Cyt (Intra)		1/330. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target

Function IGF-1 leads to the activation of AKT3, which may play a role in regulating cell survival. Capable of phosphorylating several known proteins. Truncated isoform 2/PKB gamma 1 without the second serine phosphorylation site could still be stimulated but to a lesser extent.

Tissue specificity In adult tissues, it is highly expressed in brain, lung and kidney, but weakly in heart, testis and liver. In fetal tissues, it is highly expressed in heart, liver and brain and not at all in kidney.


Sequence similarities Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. RAC subfamily. Contains 1 AGC-kinase C-terminal domain. Contains 1 PH domain. Contains 1 protein kinase domain.

Domain Binding of the PH domain to the phosphatidylinositol 3-kinase alpha (PI(3)K) results in its targeting to the plasma membrane.

Post-translational modifications Phosphorylation on Thr-305 and Ser-472 is required for full activity (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR. Ubiquitinated. When fully phosphorylated and translocated into the nucleus, undergoes 'Lys-48'-polyubiquitination catalyzed by TTC3, leading to its degradation by the proteasome.

Cellular localization Cytoplasm. Membrane. Membrane-associated after cell stimulation leading to its translocation.

Images



Western blot - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

All lanes : Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463) at 1/10000 dilution

Lane 1 : MCF7 (Human breast adenocarcinoma cell line) whole cell lysates

Lane 2 : HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysates

Lane 3 : Hep G2 (Human liver hepatocellular carcinoma) whole cell lysates

Lane 4 : A549 (Human lung carcinoma) whole cell lysates

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 56 kDa

Observed band size: 56 kDa

Blocking/Dilution buffer: 5% NFD/MTBST.




Immunocytochemistry/ Immunofluorescence - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized K562 (Human chronic myelogenous leukemia cells from bone marrow) cells labeling AKT1 + AKT2 + AKT3 with ab179463 at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/400 dilution (green). Cytoplasm and nuclear staining on K562 cell line is observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/500 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows;

1. ab179463 at 1/100 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
2. **ab7291** (anti-Tubulin mouse mAb) at 1/500 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/400 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling AKT1 + AKT2 + AKT3 with ab179463 at 1/250 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm and nucleus staining on Human renal cortex is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

All lanes : Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463) at 1/1000 dilution

Lane 1 : Human fetal brain lysate

Lane 2 : Human fetal heart lysate

Lane 3 : Human fetal kidney lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 56 kDa

Observed band size: 56 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463) at 1/1000 dilution

Lane 1 : Mouse brain lysate

Lane 2 : Mouse heart lysate

Lane 3 : Mouse kidney lysate

Lane 4 : Mouse spleen lysate

Lane 5 : Rat brain lysate

Lane 6 : Rat heart lysate

Lane 7 : Rat kidney lysate

Lane 8 : Rat spleen lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 56 kDa

Observed band size: 56 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463) at 1/10000 dilution + Xenopus muscle lysate at 10 µg


Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 56 kDa

Observed band size: 56 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

All lanes : Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463) at 1/10000 dilution

Lane 1 : AKT2 recombinant protein (HIS-tag): aa282-481

Lane 2 : AKT3 recombinant protein (HIS-tag) :aa351-479

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 56 kDa

Observed band size: 18,26 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Immunohistochemical analysis of paraffin-embedded Mouse cerebral cortex tissue labeling AKT1 + AKT2 + AKT3 with ab179463 at 1/250 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm and nucleus staining on Mouse cerebral cortex is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Immunohistochemical analysis of paraffin-embedded Rat cerebral cortex tissue labeling AKT1 + AKT2 + AKT3 with ab179463 at 1/250 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm and nucleus staining on Rat cerebral cortex is observed. Counter stained with Hematoxylin.


Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed A549 (Human lung carcinoma) cells labeling AKT1 + AKT2 + AKT3 with ab179463 at 1/330 dilution (red) compared with a rabbit monoclonal IgG isotype control (black) and a unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

AKT1 + AKT2 + AKT3 was immunoprecipitated from 1mg of MCF7 (Human breast adenocarcinoma cell line) whole cell extract with ab179463 at 1/100 dilution. Western blot was performed from the immunoprecipitate using ab179463 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution. Lane 1: MCF7 whole cell extract. Lane 2: PBS instead of MCF7 whole cell extract. Blocking and dilution buffer and concentration: 5% NFD/MBST.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-AKT1 + AKT2 + AKT3 antibody [EPR16798] (ab179463)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors